

Bone Density Regulators – Rank Ligand (RANKL) Inhibitors

Medical policy no. 30.04.45-2

Effective Date: 6/1/2025

Related medical policies:

Policy Name	Indications
Bone Density Regulators – Calcitonins	Hypercalcemia Paget's Disease Postmenopausal Osteoporosis Osteoporosis/Bone loss
Bone Density Regulators – Parathyroid Hormone Derivatives	Glucocorticoid Induced Osteoporosis Male Osteoporosis Postmenopausal Osteoporosis
Bone Density Regulators – Sclerostin Inhibitors	Postmenopausal Osteoporosis

Note: New-to-market drugs included in this class based on the Apple Health Preferred Drug List are non-preferred and subject to this prior authorization (PA) criteria. Non-preferred agents in this class require an inadequate response or documented intolerance due to severe adverse reaction or contraindication to at least TWO preferred agents. If there is only one preferred agent in the class documentation of inadequate response to ONE preferred agent is needed. If a drug within this policy receives a new indication approved by the Food and Drug Administration (FDA), medical necessity for the new indication will be determined on a case-by-case basis following FDA labeling.

To see the list of the current Apple Health Preferred Drug List (AHPDL), please visit: <https://www.hca.wa.gov/assets/billers-and-providers/apple-health-preferred-drug-list.xlsx>

Medical necessity

Drug	Medical Necessity
Denosumab (Prolia, Xgeva)	<p>Denosumab (Prolia, Xgeva) may be considered medically necessary in patients who meet the criteria described in the clinical policy below.</p> <p>If all criteria are not met, the clinical reviewer may determine there is a medically necessary need and approve on a case-by-case basis. The clinical reviewer may choose to use the reauthorization criteria when a patient has been previously established on therapy and is new to Apple Health.</p>

Clinical policy:

Clinical Criteria	
Postmenopausal Osteoporosis denosumab (Prolia)	<p>Denosumab (Prolia) may be covered when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older; AND 2. Diagnosis of osteoporosis; AND 3. Patient is a postmenopausal female; AND 4. At least ONE of the following fracture risk categories is met: <ol style="list-style-type: none"> a. Presence of fragility fractures of the hip or spine regardless of bone mineral density; OR b. T-score ≤ -2.5 in the lumbar spine, femoral neck, total hip; OR c. T-score between -1 and -2.5 with a history of recent fragility fracture of proximal humerus, pelvis, or distal forearm; OR d. T-score between -1 and -2.5 with a FRAX 10-year probability for major fracture $\geq 20\%$ or hip fracture $\geq 3\%$; AND 5. Medication will not be used in combination with other bone density regulators (e.g., bisphosphonates, raloxifene, Xgeva); AND 6. History of at least ONE of the following: <ol style="list-style-type: none"> a. Treatment with at least one Preferred Apple Health Preferred Drug List (PDL) oral or intravenous bisphosphonate medication has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 months]; OR b. Treatment with at least one Preferred Apple Health Preferred Drug List (PDL) selective estrogen receptor modulator (SERM) medication has been ineffective unless all are contraindicated, or not tolerated [minimum trial of 24 months]. <p>If ALL criteria are met, the request will be authorized for 12 months</p>
Criteria (Reauthorization)	
	<p>Denosumab (Prolia) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Criteria 5 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., patient has not suffered a fragility fracture, bone mineral density continues to improve/remain stable). <p>If ALL criteria are met, the request will be authorized for 12 months</p>
Glucocorticoid-induced osteoporosis denosumab (Prolia)	<p>Denosumab (Prolia) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older; AND

	<ol style="list-style-type: none"> 2. Diagnosis of osteoporosis; AND 3. Patient will be initiating or is continuing systemic glucocorticoid therapy at a daily dosage equivalent to ≥ 7.5 mg of prednisone and is expected to remain on glucocorticoid therapy for at least 6 months; AND 4. At least ONE of the following fracture risk categories is met: <ol style="list-style-type: none"> a. Presence of fragility fractures of the hip or spine regardless of bone mineral density; OR b. T-score ≤ -2.5 in the lumbar spine, femoral neck, total hip; OR c. T-score between -1 and -2.5 with a history of recent fragility fracture of proximal humerus, pelvis, or distal forearm; OR d. T-score between -1 and -2.5 with a FRAX 10-year probability for major fracture $\geq 20\%$ or hip fracture $\geq 3\%$; AND 5. Medication will not be used in combination with other bone density regulators (e.g., bisphosphonates, raloxifene, Xgeva); AND 6. Treatment with at least one Preferred Apple Health Preferred Drug List (PDL) oral or intravenous bisphosphonate medication indicated for glucocorticoid induced osteoporosis has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 months]. <p>If ALL criteria are met, the request will be authorized for 12 months</p> <p>Criteria (Reauthorization)</p> <p>Denosumab (Prolia) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient has not suffered a fragility fracture while on treatment; AND 2. Criteria 5 above continues to be met; AND 3. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., patient has not suffered a fragility fracture, bone mineral density continues to improve/remain stable). <p>If ALL criteria are met, the request will be authorized for 12 months</p>
Treatment of bone loss in men with prostate cancer denosumab (Prolia)	<p>Denosumab (Prolia) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older; AND 2. Patient has a diagnosis of bone loss or osteoporosis indicated by one or more of the following: <ol style="list-style-type: none"> a. Presence of fragility fractures of the hip or spine regardless of bone mineral density; OR b. T-score ≤ -2.5 in the lumbar spine, femoral neck, total hip; OR c. T-score between -1 and -2.5 with a history of recent fragility fracture of proximal humerus, pelvis, or distal forearm; OR

	<p>d. T-score between -1 and -2.5 with a FRAX 10-year probability for major fracture $\geq 20\%$ or hip fracture $\geq 3\%$; AND</p> <p>3. Patient is currently receiving androgen deprivation therapy (ADT) (e.g., leuprolide, degarelix, relugolix) for non-metastatic prostate cancer unless all are contraindicated or not tolerated; AND</p> <p>4. Medication will not be used in combination with denosumab (Xgeva); AND</p> <p>5. Treatment with at least one Preferred Apple Health Preferred Drug List (PDL) oral or intravenous bisphosphonate medication has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 months].</p> <p>If ALL criteria are met, the request will be authorized for 12 months</p> <p>Criteria (Reauthorization)</p> <p>Denosumab (Prolia) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Criteria 3 and 4 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., patient has not suffered a fragility fracture, bone mineral density continues to improve/remains stable). <p>If ALL criteria are met, the request will be authorized for 12 months</p>
Treatment of bone loss in women with breast cancer denosumab (Prolia)	<p>Denosumab (Prolia) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older; AND 2. Patient has a diagnosis of bone loss or osteoporosis indicated by one or more of the following: <ol style="list-style-type: none"> a. Presence of fragility fractures of the hip or spine regardless of bone mineral density; OR b. T-score ≤ -2.5 in the lumbar spine, femoral neck, total hip; OR c. T-score between -1 and -2.5 with a history of recent fragility fracture of proximal humerus, pelvis, or distal forearm; OR d. T-score between -1 and -2.5 with a FRAX 10-year probability for major fracture $\geq 20\%$ or hip fracture $\geq 3\%$; AND 3. Patient is receiving adjuvant aromatase inhibitor therapy (e.g., anastrozole, exemestane, letrozole) for breast cancer unless all are contraindicated or not tolerated; AND 4. Medication will not be used in combination with denosumab (Xgeva); AND 5. Treatment with at least one Preferred Apple Health Preferred Drug List (PDL) oral or intravenous bisphosphonate medication has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 months].

	<p>If ALL criteria are met, the request will be authorized for 12 months</p> <p>Criteria (Reauthorization)</p> <p>Denosumab (Prolia) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Criteria 4 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., patient has not suffered a fragility fracture, bone mineral density continues to improve/remain stable). <p>If ALL criteria are met, the request will be authorized for 12 months</p>
<p>Multiple Myeloma and bone metastasis from solid tumors denosumab (Xgeva)</p>	<p>Denosumab (Xgeva) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older; AND 2. Patient has one of the following: <ol style="list-style-type: none"> a. Diagnosis of multiple myeloma with skeletal-related events (i.e., radiation to bone, pathologic fracture, surgery to bone, and spinal cord compression); OR b. Bone metastases from solid tumors (i.e., metastatic breast cancer, metastatic castration-resistant prostate cancer, or metastatic lung cancer) AND 3. History of failure, contraindication, or intolerance to zoledronic acid; AND 4. Medication will not be used in combination with denosumab (Prolia) <p>If ALL criteria are met, the request will be authorized for 12 months</p> <p>Criteria (Reauthorization)</p> <p>Denosumab (Xgeva) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Criteria 4 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., patient has not suffered a fracture, bone mineral density continues to improve/remain stable). <p>If ALL criteria are met, the request will be authorized for 12 months</p>
<p>Giant cell tumor of bone denosumab (Xgeva)</p>	<p>Denosumab (Xgeva) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 12 years of age or older AND skeletally mature; AND 2. Diagnosis of giant cell tumor of the bone; AND <ol style="list-style-type: none"> a. Disease is unresectable or surgical resection is likely to result in severe morbidity; OR b. Disease is recurrent or metastatic 3. Medication will not be used in combination with denosumab (Prolia).

	<p>If ALL criteria are met, the request will be authorized for 12 months</p> <p>Criteria (Reauthorization)</p> <p>Denosumab (Xgeva) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Criteria 3 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., decrease in tumor size or spread of tumor). <p>If ALL criteria are met, the request will be authorized for 12 months</p>
Hypercalcemia of malignancy denosumab (Xgeva)	<p>Denosumab (Xgeva) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older; AND 2. Diagnosis of hypercalcemia of malignancy; AND 3. Baseline corrected serum calcium > 12.5 mg/dL 3. Treatment with at least one Preferred Apple Health Preferred Drug List (PDL) oral or intravenous bisphosphonate medication has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 months]. 4. Medication will not be used in combination with denosumab (Prolia). <p>If ALL criteria are met, the request will be authorized for 12 months</p> <p>Criteria (Reauthorization)</p> <p>Denosumab (Xgeva) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Criteria 4 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., decrease in corrected serum calcium). <p>If ALL criteria are met, the request will be authorized for 12 months</p>

Dosage and quantity limits

Drug	Indication	FDA Approved Dosing	Dosage Form and Quantity Limit
denosumab (Prolia)	<p>Glucocorticoid-induced osteoporosis</p> <p>Postmenopausal Osteoporosis</p> <p>Treatment of bone loss in women with breast cancer</p>	60 mg every 6 months	<ul style="list-style-type: none"> 60 mg/1 mL prefilled syringe 1 mL/168 days

	Treatment of bone loss in men with prostate cancer		
denosumab (Xgeva)	Giant cell tumor of bone	120 mg every 4 weeks with additional 120 mg doses on Days 8 and 15 of the first month of therapy	<ul style="list-style-type: none"> • 120 mg/1.7 mL vial: <ul style="list-style-type: none"> ○ Loading: 5.1 mL/28 days ○ Maintenance: 1.7 mL/28 days
denosumab (Xgeva)	Multiple Myeloma and bone metastasis from solid tumors	120 mg every 4 weeks	<ul style="list-style-type: none"> • 120 mg/1.7 mL vial: 1.7 mL/28 days

Coding:

HCPSC Code	Description
J0897	Injection, denosumab, 1 mg

Background:

Osteoporosis is characterized by decreased bone mass and increased fracture risk, most commonly at the spine, hip, and wrist. The definition of osteoporosis with high risk of fracture is defined for men and women as BMD T-score of spine, femoral neck, and/or total hip <-2.5 without fracture, having history of hip or vertebral fracture regardless of BMD, T-score ≤ -1 and a history of recent fracture of proximal humerus, pelvis, or distal forearm, T-score between -1.0 and -2.5 in the spine, femoral neck, or total hip with a -20% 10-year FRAX risk of any fracture or -3% risk of hip fracture, and receiving long-term glucocorticoid doses greater than or equal to prednisone 7.5mg per day. The treatment of osteoporosis consists of lifestyle management (e.g., adequate calcium and vitamin D, exercise, smoking cessation, fall prevention measures, and avoidance of heavy alcohol use) and pharmacologic therapy. Patients with the highest risk of fracture are expected to derive the greatest benefit from medication therapy. The 2020 AACE/ACE treatment guideline recommendations are as follows:

1. Initial treatment for high fracture risk: alendronate, denosumab, risedronate, or zoledronic acid
2. Treatment for very-high fracture risk or patients who cannot tolerate or adhere to oral bisphosphonates: zoledronic acid, abaloparatide, denosumab, romosozumab, teriparatide

Additionally, the 2020 Endocrine Society guidelines recommend bisphosphonates as initial treatment for high-risk patients, while denosumab may be considered as an alternative initial treatment. For patients with a very high risk of fracture, teriparatide and abaloparatide are recommended. It is recommended that antiresorptive therapies follow treatment with parathyroid hormones.

References

1. Prolia [package insert]. Thousand Oaks, CA; Amgen, Inc.; January 2023.
2. Xgeva [package insert]. Thousand Oaks, CA; Amgen, Inc.; June 2020.
3. Camacho PM, Petak SM, Binkley N, et al. American Association of Clinical Endocrinologists/American College of Endocrinology clinical practice guidelines for the diagnosis and treatment of postmenopausal osteoporosis-2020 update. Endocr Pract. 2020;26(Suppl 1):1-46.

4. Shoback D, Rosen CJ, Black DM, Cheung AM, Murad MH, Eastell R. Pharmacological management of osteoporosis in postmenopausal women: an Endocrine Society guideline update. J Clin Endocrinol Metab. 2020;105(3):dgaa048.

History

Approved Date	Effective Date	Version	Action and Summary of Changes
10/16/2025	06/01/2025	30.04.45-2	<ul style="list-style-type: none"> Removed “indicated for glucocorticoid induced osteoporosis” from clinical criteria for hypercalcemia of malignancy
10/16/2024	06/01/2025	30.04.45-2	<p>Approved by DUR Board</p> <ul style="list-style-type: none"> Removed denosumab (Prolia) from broader Bone Density Regulators policy and aged denosumab (Xgeva) to the RankL policy Broke out policy criteria by indication Updated verbiage around prior treatment with bisphosphonates Added language in around requirement to be high risk of fracture Added criteria for denosumab (Xgeva) for respective indications
	10/01/2019	30.04.00-1	<ul style="list-style-type: none"> 07.31.2019 Updated abaloparatide, teriparatide criteria 05.31.2019 Updated abaloparatide, teriparatide, and denosumab reauthorization criteria 04.01.2019 Added Brands with Generic Equivalents policy; updated abaloparatide and teriparatide clinical policies 04.18.2018 New Policy