

Bone Density Regulators – Parathyroid Hormone Analogs

Medical policy no. 30.04.40-2

Effective Date: 6/1/2025

Related medical policies:

Policy Name	Indications
Bone Density Regulators – Calcitonins	Hypercalcemia Paget's Disease Postmenopausal Osteoporosis Osteoporosis/Bone loss
Bone Density Regulators- RANKL Inhibitors	Giant cell tumor of bone Glucocorticoid-induced Osteoporosis Hypercalcemia of malignancy Multiple Myeloma and bone metastasis from solid tumors Postmenopausal Osteoporosis Treatment of bone loss in men prostate cancer Treatment of bone loss in women with breast cancer
Bone Density Regulators – Sclerostin Inhibitors	Postmenopausal Osteoporosis

Note: New-to-market drugs included in this class based on the Apple Health Preferred Drug List are non-preferred and subject to this prior authorization (PA) criteria. Non-preferred agents in this class require an inadequate response or documented intolerance due to severe adverse reaction or contraindication to at least TWO preferred agents. If there is only one preferred agent in the class documentation of inadequate response to ONE preferred agent is needed. If a drug within this policy receives a new indication approved by the Food and Drug Administration (FDA), medical necessity for the new indication will be determined on a case-by-case basis following FDA labeling.

To see the list of the current Apple Health Preferred Drug List (AHPDL), please visit: <https://www.hca.wa.gov/assets/billers-and-providers/apple-health-preferred-drug-list.xlsx>

Medical necessity

Drug	Medical Necessity
abaloparatide (Tymlos) teriparatide (Forteo)	<p>Abaloparatide (Tymlos) and teriparatide (Forteo) may be considered medically necessary in patients who meet the criteria described in the clinical policy below.</p> <ul style="list-style-type: none"> Non-preferred brand name products on the Apple Health Preferred Drug List with an A-rated generic equivalent must also meet criteria in Non-Clinical Policy No. 0001 (NC-001). <p>If all criteria are not met, the clinical reviewer may determine there is a medically necessary need and approve on a case-by-case basis. The clinical reviewer may choose to use the reauthorization criteria when a</p>

patient has been previously established on therapy and is new to Apple Health.

Clinical policy:

Clinical Criteria	
Postmenopausal Osteoporosis	<p>Abaloparatide (Tymlos) and teriparatide (Forteo) may be covered when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older; AND 2. Diagnosis of osteoporosis; AND 3. Patient is a postmenopausal female; AND 4. At least ONE of the following fracture risk categories is met: <ol style="list-style-type: none"> a. Presence of fragility fractures of the hip or spine regardless of bone mineral density; OR b. T-score ≤ -2.5 in the lumbar spine, femoral neck, total hip; OR c. T-score between -1 and -2.5 with a history of recent fragility fracture of proximal humerus, pelvis, or distal forearm; OR d. T-score between -1 and -2.5 with a FRAX 10-year probability for major fracture $\geq 20\%$ or hip fracture $\geq 3\%$; AND 5. Treatment duration has not exceeded a total of 24 months of cumulative use of a parathyroid hormone during their lifetime; OR 6. <u>For teriparatide only:</u> Patient received treatment with a parathyroid hormone for more than 24 months during their lifetime; AND <ol style="list-style-type: none"> a. Patient remains, or has returned to, having high or very high fracture risk (e.g., a fracture in the past 12 months, a fracture while on osteoporosis therapy, a history of multiple fractures, fractures while on long-term glucocorticoids, T-score ≤ -3.0, high risk for falls or a history of injurious falls, a FRAX 10-year probability for major fracture $>30\%$ or hip fracture $>4.5\%$, etc.); AND 7. Medication will not be used in combination with other bone density regulators (e.g., bisphosphonates, raloxifene, RANKL inhibitor); AND 8. History of at least ONE of the following: <ol style="list-style-type: none"> a. Minimum trial of 12 months with at least one Preferred Apple Health Preferred Drug List (PDL) oral or intravenous bisphosphonate medication has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 months]; OR b. Minimum trial of 24 months with at least one Preferred Apple Health Preferred Drug List (PDL) selective estrogen receptor modulator (SERM) medication has been ineffective unless all are contraindicated, or not tolerated [minimum trial of 24 months]; OR

	<ul style="list-style-type: none"> c. Treatment with denosumab has been ineffective unless contraindicated or not tolerated [minimum trial of 12 months]; OR d. Abaloparatide or teriparatide may be considered medically necessary for first line therapy for severe osteoporosis defined as either of the following: <ul style="list-style-type: none"> i. T-score \leq -2.5 with a fragility fracture; OR ii. History of multiple fragility fractures; OR iii. T-score \leq -3 regardless of previous therapy. <p>If ALL criteria are met, the request will be authorized for up to 12 months, unless total combined duration of parathyroid hormone analog would exceed 2 years.</p> <p>Criteria (Reauthorization)</p> <p>Abaloparatide (Tymlos) and teriparatide (Forteo) may be approved when all the following documented criteria are met:</p> <ul style="list-style-type: none"> 1. Criteria 7 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., patient has not suffered a fragility fracture, bone mineral density continues to improve/remains stable). <p><u>For abaloparatide:</u> If all the criteria are met, approve for up to 12 months, unless total combined duration of parathyroid hormone analog would exceed 2 years.</p> <p><u>For teriparatide:</u> If all criteria are met, approve for up to 12 months.</p>
Male Osteoporosis	<p>Abaloparatide (Tymlos) and teriparatide (Forteo) may be covered when all the following documented criteria are met:</p> <ul style="list-style-type: none"> 1. Patient is 18 years of age or older; AND 2. Diagnosis of osteoporosis; AND 3. Patient is a biological male; AND 4. At least ONE of the following fracture risk categories is met: <ul style="list-style-type: none"> a. Presence of fragility fractures of the hip or spine regardless of bone mineral density; OR b. T-score \leq -2.5 in the lumbar spine, femoral neck, total hip; OR c. T-score between -1 and -2.5 with a history of recent fragility fracture of proximal humerus, pelvis, or distal forearm; OR d. T-score between -1 and -2.5 with a FRAX 10-year probability for major fracture \geq20% or hip fracture \geq3%; AND 5. Treatment duration has not exceeded a total of 24 months of cumulative use of a parathyroid hormone during their lifetime; OR

	<p>6. <u>For teriparatide only</u>: Patient received treatment with a parathyroid hormone for more than 24 months during their lifetime; AND</p> <p>a. Patient remains, or has returned to, having high or very high fracture risk (e.g., a fracture in the past 12 months, a fracture while on osteoporosis therapy, a history of multiple fractures, fractures while on long-term glucocorticoids, T-score ≤ -3.0, high risk for falls or a history of injurious falls, a FRAX 10-year probability for major fracture $>30\%$ or hip fracture $>4.5\%$, etc.); AND</p> <p>7. Medication will not be used in combination with other bone density regulators (e.g., bisphosphonates, raloxifene, RANKL inhibitor); AND</p> <p>8. Treatment with at least one Preferred Apple Health Preferred Drug List (PDL) oral or intravenous bisphosphonate medication indicated for male osteoporosis has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 months]; OR</p> <p>9. Abaloparatide or teriparatide may be considered medically necessary for first line therapy for severe osteoporosis defined as either of the following:</p> <ul style="list-style-type: none"> i. T-score ≤ -2.5 with a fragility fracture; OR ii. History of multiple fragility fractures; OR iii. T-score ≤ -3 regardless of previous therapy. <p>If ALL criteria are met, the request will be authorized for up to 12 months, unless total combined duration of parathyroid hormone analog would exceed 2 years.</p> <p>Criteria (Reauthorization)</p> <p>Abaloparatide (Tymlos) and teriparatide (Forteo) may be approved when all the following documented criteria are met:</p> <ul style="list-style-type: none"> 1. Criteria 7 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., patient has not suffered a fragility fracture, bone mineral density continues to improve/remains stable). <p><u>For abaloparatide</u>: If all the criteria are met, approve for up to 12 months, unless total combined duration of parathyroid hormone analog would exceed 2 years.</p> <p><u>For teriparatide</u>: If all criteria are met, approve for up to 12 months.</p>
Glucocorticoid Induced Osteoporosis	<p>Teriparatide (Forteo) may be covered when all the following documented criteria are met:</p> <ul style="list-style-type: none"> 1. Patient is 18 years of age or older; AND 2. Diagnosis of osteoporosis; AND 3. Patient has a history of or is currently taking sustained systemic glucocorticoid therapy (daily dosage equivalent to ≥ 5 mg of prednisone) [minimum use of 3 months]; AND

	<ol style="list-style-type: none"> 4. At least ONE of the following fracture risk categories is met: <ol style="list-style-type: none"> a. Presence of fragility fractures of the hip or spine regardless of bone mineral density; OR b. T-score ≤ -2.5 in the lumbar spine, femoral neck, total hip; OR c. T-score between -1 and -2.5 with a history of recent fragility fracture of proximal humerus, pelvis, or distal forearm; OR d. T-score between -1 and -2.5 with a FRAX 10-year probability for major fracture $\geq 20\%$ or hip fracture $\geq 3\%$; AND 5. Treatment duration has not exceeded a total of 24 months of cumulative use of a parathyroid hormone during their lifetime; OR 6. Patient received treatment with a parathyroid hormone for more than 24 months during their lifetime; AND <ol style="list-style-type: none"> a. Patient remains, or has returned to, having high or very high fracture risk (e.g., a fracture in the past 12 months, a fracture while on osteoporosis therapy, a history of multiple fractures, fractures while on long-term glucocorticoids, T-score ≤ -3.0, high risk for falls or a history of injurious falls, a FRAX 10-year probably for 7. Medication will not be used in combination with other bone density regulators (e.g., bisphosphonates, raloxifene, RANKL inhibitor); AND 8. Treatment with at least one Preferred Apple Health Preferred Drug List (PDL) oral or intravenous bisphosphonate medication indicated for glucocorticoid induced osteoporosis has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 months]; OR 9. Teriparatide may be considered medically necessary for first line therapy for severe osteoporosis defined as either of the following: <ol style="list-style-type: none"> i. T-score ≤ -2.5 with a fragility fracture; OR ii. History of multiple fragility fractures; OR iii. T-score ≤ -3 regardless of previous therapy. <p>If ALL criteria are met, the request will be authorized for up to 12 months, unless total combined duration of parathyroid hormone analog would exceed 2 years.</p>
	<p>Criteria (Reauthorization)</p> <p>Teriparatide (Forteo) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Criteria 7 above continues to be met; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., patient has not suffered a fragility fracture, bone mineral density continues to improve/remain stable). <p>If all the criteria are met, approve for up to 12 months.</p>

Dosage and quantity limits

Drug	Indication	FDA Approved Dosing	Dosage Form and Quantity Limit
Tymlos (abaloparatide)	Postmenopausal Osteoporosis	80 mcg subQ once daily	3120 mcg/1.56 mL prefilled pen 1.56 mL/28 days
	Male Osteoporosis		
Forteo	Postmenopausal Osteoporosis	20 mcg subQ once daily	600 mcg/2.4 mL prefilled pen 2.4 mL/28 days
	Male Osteoporosis		
	Glucocorticoid Induced Osteoporosis		
Teriparatide (generic)	Postmenopausal Osteoporosis	20 mcg subQ once daily	600 mcg/2.4 mL prefilled pen 2.4 mL/28 days
	Male Osteoporosis		
	Glucocorticoid Induced Osteoporosis		

Coding:

HCPSC Code	Description
J3110	Injection, teriparatide, 10 mcg

Background:

Osteoporosis is characterized by decreased bone mass and increased fracture risk, most commonly at the spine, hip, and wrist. The definition of osteoporosis with high risk of fracture is defined for men and women as BMD T-score of spine, femoral neck, and/or total hip <-2.5 without fracture, having history of hip or vertebral fracture regardless of BMD, T-score ≤ -1 and a history of recent fracture of proximal humerus, pelvis, or distal forearm, T-score between -1.0 and -2.5 in the spine, femoral neck, or total hip with a -20% 10-year FRAX risk of any fracture or -3% risk of hip fracture, and receiving long-term glucocorticoid doses greater than or equal to prednisone 7.5mg per day. The treatment of osteoporosis consists of lifestyle management (e.g., adequate calcium and vitamin D, exercise, smoking cessation, fall prevention measures, and avoidance of heavy alcohol use) and pharmacologic therapy. Patients with the highest risk of fracture are expected to derive the greatest benefit from medication therapy. The 2020 AACE/ACE treatment guideline recommendations are as follows:

1. Initial treatment for high fracture risk: alendronate, denosumab, risedronate, or zoledronic acid
2. Treatment for very-high fracture risk or patients who cannot tolerate or adhere to oral bisphosphonates: zoledronic acid, abaloparatide, denosumab, romosozumab, teriparatide

Additionally, the 2020 Endocrine Society guidelines recommend bisphosphonates as initial treatment for high-risk patients, while denosumab may be considered as an alternative initial treatment. For patients with a very high risk of fracture, teriparatide and abaloparatide are recommended. It is recommended that antiresorptive therapies follow treatment with parathyroid hormones.

References

1. Prolia [package insert]. Thousand Oaks, CA; Amgen, Inc.; January 2023.
2. Xgeva [package insert]. Thousand Oaks, CA; Amgen, Inc.; June 2020.
3. Camacho PM, Petak SM, Binkley N, et al. American Association of Clinical Endocrinologists/American College of Endocrinology clinical practice guidelines for the diagnosis and treatment of postmenopausal osteoporosis-2020 update. Endocr Pract. 2020;26(Suppl 1):1-46.
4. Shoback D, Rosen CJ, Black DM, Cheung AM, Murad MH, Eastell R. Pharmacological management of osteoporosis in postmenopausal women: an Endocrine Society guideline update. J Clin Endocrinol Metab. 2020;105(3):dgaa048.

History

Approved Date	Effective Date	Version	Action and Summary of Changes
10/16/2024	06/01/2025	30.04.40-2	<p>Approved by DUR Board</p> <ul style="list-style-type: none"> - Split out Bone Density Regulator policy into different policies
	10/01/2019	30.04.00-1	<ul style="list-style-type: none"> • 07.31.2019 Updated abaloparatide, teriparatide criteria • 05.31.2019 Updated abaloparatide, teriparatide, and denosumab reauthorization criteria • 04.01.2019 Added Brands with Generic Equivalents policy; updated abaloparatide and teriparatide clinical policies • 04.18.2018 New Policy