Appendix C: MCO Comparison Results

Appendix C contains a subset of the information included in Appendix B for all the performance measures by MCO and by region and is available publicly.

Comagine Health C-1

Legend:

A

Measure result is statistically significant above prior year (p < 0.05)

•

Measure result is statistically significant below prior year (p < 0.05)



Can report benchmark Rotated measures*

NR There was no data to report.

| Measure Description | MCO | MY 2018 Rate | MY 2019 Rate | MY 2020 Rate | MY 2020 Performance** |
|--|------------|--------------|--------------|--------------|-----------------------|
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 72.8% | 73.9% | 72.8% | Below 50th |
| Activity for Children/Adolescents (WCC), BMI Percentile, 3-11 | AMG | 73.4% | 73.4% | 69.8% | Below 50th |
| Years | CCW | 64.3% | 66.3% | 64.5% | Below 50th |
| | CHPW | 79.4% | 76.4% | 75.3% | At 50th |
| | MHW | 76.6% | 76.6% | 75.9% | At 50th |
| | UHC | 54.9% | 64.9% | 66.5% | Below 50th |
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 71.2% | 71.7% | 64.5% | Below 50th |
| Activity for Children/Adolescents (WCC), BMI Percentile, 12-17 | AMG | 71.3% | 71.3% | 69.2% | At 50th |
| Years | CCW | 52.1% | 60.0% | 57.4% | Below 50th |
| | CHPW | 69.5% | 78.3% | 72.6% | At 50th |
| | MHW | 79.7% | 73.7% | 64.6% | Below 50th |
| | UHC | 55.8% | 63.7% | 57.2% | Below 50th |
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 72.2% | 73.1% | 69.6% | Below 50th |
| Activity for Children/Adolescents (WCC), BMI Percentile, Total | AMG | 72.7% | 72.7% | 69.6% | Below 50th |
| | CCW | 60.1% | 64.0% | 61.8% | Below 50th |
| | CHPW | 75.7% | 77.1% | 74.2% | At 50th |
| | MHW | 77.6% | 75.7% | 71.5% | Below 50th |

[†] Statewide Weighted Rate

^{*} In March 2020, NCQA recognized that COVID-19 would likely impact plans' ability to collect medical record data due to travel bans, quarantines, and efforts to minimize risk to staff. Therefore, NCQA allowed Medicaid plans participating in HEDIS reporting the option of submitting 2019 rates for their hybrid measures ("rotated measures").

^{** 2021} performance indicates whether a measure is statistically different than national benchmarks.

^{***} Rates suppressed when the denominator is less than 30.

| | UHC | 55.2% | 64.5% | 63.3% | Below 50th |
|--|------------|-------|-------|---------|------------|
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 62.8% | 64.8% | 64.8% | Below 50th |
| Activity for Children/Adolescents (WCC), Nutrition Counseling, 3- | AMG | 62.8% | 62.8% | 58.1% | Below 50th |
| 11 Years | CCW | 70.3% | 76.6% | 70.7% | At 50th |
| | CHPW | 69.5% | 64.1% | 70.1% | At 50th |
| | MHW | 60.1% | 62.8% | 62.9% | Below 50th |
| | UHC | 56.0% | 60.5% | 67.3% | At 50th |
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 60.0% | 59.2% | 51.4% | Below 50th |
| Activity for Children/Adolescents (WCC), Nutrition Counseling, 12- | - AMG | 55.8% | 55.8% | 54.8% | Below 50th |
| 17 Years | CCW | 62.0% | 59.3% | 61.9% | At 50th |
| | CHPW | 64.9% | 60.8% | 59.2% | Below 50th |
| | MHW | 58.6% | 59.1% | 46.2% | Below 50th |
| | UHC | 57.2% | 57.8% | 53.1% | Below 50th |
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 61.8% | 62.8% | 59.7% | Below 50th |
| Activity for Children/Adolescents (WCC), Nutrition Counseling, | AMG | 60.6% | 60.6% | 56.9% | Below 50th |
| Total | CCW | 67.4% | 70.3% | 67.4% | At 50th |
| | CHPW | 67.8% | 62.9% | 65.7% | At 50th |
| | MHW | 59.6% | 61.6% | 56.5% | Below 50th |
| | UHC | 56.4% | 59.6% | 62.3% | Below 50th |
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 55.3% | 58.0% | 58.7% | Below 50th |
| Activity for Children/Adolescents (WCC), Physical Activity | AMG | 52.8% | 52.8% | 53.6% | Below 50th |
| Counseling, 3-11 Years | CCW | 64.3% | 69.3% | 63.3% | At 50th |
| | CHPW | 66.4% | 61.2% | 60.2% | Below 50th |
| | MHW | 52.2% | 56.2% | 57.3% | Below 50th |
| | UHC | 43.2% | 48.9% | 62.8% 🔺 | At 50th |
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 61.9% | 59.9% | 52.4% | Below 50th |
| Activity for Children/Adolescents (WCC), Physical Activity | AMG | 57.4% | 57.4% | 54.8% | Below 50th |
| Counseling, 12-17 Years | CCW | 65.5% | 62.0% | 61.9% | At 50th |
| | CHPW | 68.7% | 67.1% | 58.6% | Below 50th |
| | MHW | 59.4% | 56.9% | 48.1% | Below 50th |
| | UHC | 60.1% | 61.5% | 53.8% | Below 50th |
| Weight Assessment and Counseling for Nutrition and Physical | Statewide† | 57.5% | 58.6% | 56.3% | Below 50th |
| Activity for Children/Adolescents (WCC), Physical Activity | AMG | 54.3% | 54.3% | 54.0% | Below 50th |
| Counseling, Total | CCW | 64.7% | 66.7% | 62.8% | At 50th |

| | CHPW | 67.2% | 63.4% | 59.5% Below 50th |
|--|------------|-------|-------|---------------------------|
| | MHW | 54.5% | 56.4% | 53.8% Below 50th |
| | UHC | 48.9% | 53.0% | 59.6% Below 50th |
| Childhood Immunization Status (CIS), DTaP | Statewide† | 76.9% | 77.7% | 72.8% ▼ Below 50th |
| | AMG | 73.5% | 75.2% | 75.2% At 50th |
| | CCW | 78.8% | 84.9% | 81.5% At 75th |
| | CHPW | 76.9% | 76.4% | 78.8% At 75th |
| | MHW | 76.6% | 76.6% | 69.4% ▼ Below 50th |
| | UHC | 77.6% | 77.6% | 70.1% Below 50th |
| Childhood Immunization Status (CIS), IPV | Statewide† | 88.9% | 89.2% | 86.1% ▼ Below 50th |
| | AMG | 87.8% | 87.8% | 88.8% At 50th |
| | CCW | 91.7% | 94.2% | 92.7% At 75th |
| | CHPW | 89.1% | 89.3% | 92.0% At 75th |
| | MHW | 88.1% | 88.1% | 82.4% ▼ Below 50th |
| | UHC | 89.5% | 89.5% | 87.1% At 50th |
| Childhood Immunization Status (CIS), MMR | Statewide† | 87.3% | 87.8% | 85.3% ▼ Below 50th |
| | AMG | 85.6% | 87.6% | 88.1% At 50th |
| | CCW | 91.0% | 94.2% | 89.5% At 50th |
| | CHPW | 88.1% | 88.6% | 89.1% At 50th |
| | MHW | 85.6% | 85.6% | 82.6% Below 50th |
| | UHC | 89.3% | 89.3% | 86.4% At 50th |
| Childhood Immunization Status (CIS), HIB | Statewide† | 88.2% | 88.4% | 85.2% ▼ Below 50th |
| | AMG | 86.6% | 86.6% | 87.6% At 50th |
| | CCW | 91.0% | 92.9% | 92.2% At 75th |
| | CHPW | 87.8% | 88.6% | 90.8% At 75th |
| | MHW | 87.6% | 87.6% | 81.6% ▼ Below 50th |
| | UHC | 88.3% | 88.3% | 85.9% At 50th |
| Childhood Immunization Status (CIS), Hepatitis B | Statewide† | 88.3% | 88.0% | 84.5% ▼ Below 50th |
| | AMG | 88.3% | 86.6% | 89.5% At 50th |
| | CCW | 92.7% | 92.9% | 92.7% At 75th |
| | CHPW | 90.3% | 89.3% | 91.5% At 50th |
| | MHW | 86.6% | 86.6% | 79.4% ▼ Below 50th |
| | UHC | 87.8% | 87.8% | 86.4% At 50th |
| Childhood Immunization Status (CIS), VZV | Statewide† | 86.5% | 87.1% | 84.1% ▼ Below 50th |

| 1 | | | | |
|---|------------|-------|-------|---------------------------|
| | AMG | 84.2% | 85.9% | 86.6% At 50th |
| | CCW | 89.8% | 93.2% | 87.8% At 50th |
| | CHPW | 87.6% | 88.6% | 88.6% At 50th |
| | MHW | 85.2% | 85.2% | 81.3% Below 50th |
| | UHC | 88.3% | 88.3% | 85.6% At 50th |
| Childhood Immunization Status (CIS), Pneumococcal | Statewide† | 76.8% | 77.6% | 72.9% T Below 50th |
| | AMG | 74.5% | 75.7% | 75.4% At 50th |
| | CCW | 80.3% | 84.7% | 80.8% At 75th |
| | CHPW | 76.4% | 78.1% | 78.1% At 50th |
| | MHW | 75.7% | 75.7% | 69.3% ▼ Below 50th |
| | UHC | 79.3% | 79.3% | 72.0% Below 50th |
| Childhood Immunization Status (CIS), Hepatitis A | Statewide† | 83.1% | 83.7% | 81.1% ▼ Below 50th |
| | AMG | 82.0% | 83.7% | 85.4% At 50th |
| | CCW | 88.1% | 91.5% | 88.6% At 75th |
| | CHPW | 84.4% | 85.9% | 85.9% At 50th |
| | MHW | 80.8% | 80.8% | 77.2% Below 50th |
| | UHC | 85.4% | 85.4% | 81.3% At 50th |
| Childhood Immunization Status (CIS), Rotavirus | Statewide† | 68.5% | 69.3% | 69.8% ▼ Below 50th |
| | AMG | 69.8% | 71.5% | 70.3% At 50th |
| | CCW | 72.5% | 79.3% | 76.4% At 50th |
| | CHPW | 72.7% | 72.0% | 71.5% At 50th |
| | MHW | 65.0% | 65.0% | 67.3% Below 50th |
| | UHC | 72.5% | 72.5% | 72.0% At 50th |
| Childhood Immunization Status (CIS), Influenza | Statewide† | 55.4% | 55.3% | 55.0% ▼ At 50th |
| | AMG | 53.3% | 54.7% | 52.8% At 50th |
| | CCW | 60.6% | 67.4% | 69.1% Above 75th |
| | CHPW | 62.3% | 56.0% | 61.1% At 75th |
| | MHW | 52.1% | 52.1% | 50.1% At 50th |
| | UHC | 55.7% | 55.7% | 56.9% At 75th |
| Childhood Immunization Status (CIS), Combo 2 | Statewide† | 73.2% | 74.0% | 68.3% ▼ Below 50th |
| | AMG | 69.8% | 71.5% | 72.3% At 50th |
| | CCW | 77.1% | 82.2% | 77.9% At 75th |
| | CHPW | 74.2% | 74.7% | 77.1% At 75th |
| | MHW | 72.0% | 72.0% | 63.4% ▼ Below 50th |
| 1 | | | | |

| | UHC | 74.5% 74.5% | 66.9% At 50th |
|--|------------|-------------|--------------------|
| Childhood Immunization Status (CIS), Combo 3 | Statewide† | 70.0% 70.7% | 64.8% ▼ Below 50th |
| | AMG | 66.7% 68.9% | 68.1% At 50th |
| | CCW | 75.2% 78.8% | 73.5% At 75th |
| | CHPW | 70.6% 72.0% | 72.8% At 75th |
| | MHW | 68.6% 68.6% | 60.2% ▼ Below 50th |
| | UHC | 71.0% 71.0% | 64.0% At 50th |
| hildhood Immunization Status (CIS), Combo 4 | Statewide† | 67.8% 68.9% | 63.9% ▼ Below 50th |
| | AMG | 64.7% 68.1% | 67.4% At 50th |
| | CCW | 73.7% 78.6% | 73.5% At 75th |
| | CHPW | 69.3% 71.5% | 71.5% At 75th |
| | MHW | 65.7% 65.7% | 59.1% ▼ Below 50th |
| | UHC | 69.6% 69.6% | 63.3% At 50th |
| Childhood Immunization Status (CIS), Combo 5 | Statewide† | 59.6% 60.7% | 57.7% ▼ Below 50th |
| | AMG | 58.4% 62.0% | 59.4% At 50th |
| | CCW | 63.5% 70.1% | 65.0% At 75th |
| | CHPW | 62.8% 63.7% | 64.0% At 50th |
| | MHW | 56.7% 56.7% | 54.1% Below 50th |
| | UHC | 63.5% 63.5% | 57.9% At 50th |
| Childhood Immunization Status (CIS), Combo 6 | Statewide† | 47.7% 47.8% | 46.1% ▼ At 50th |
| | AMG | 44.5% 48.7% | 45.7% At 50th |
| | CCW | 54.0% 60.1% | 58.9% Above 75th |
| | CHPW | 53.5% 47.7% | 53.0% At 75th |
| | MHW | 44.5% 44.5% | 41.2% Below 50th |
| | UHC | 48.4% 48.4% | 47.5% At 75th |
| hildhood Immunization Status (CIS), Combo 7 | Statewide† | 58.3% 59.7% | 57.0% ▼ Below 50th |
| | AMG | 57.2% 61.8% | 58.6% At 50th |
| | CCW | 62.5% 69.8% | 65.0% At 75th |
| | CHPW | 61.6% 63.3% | 62.8% At 50th |
| | MHW | 55.2% 55.2% | 53.3% Below 50th |
| | UHC | 62.3% 62.3% | 57.2% At 50th |
| hildhood Immunization Status (CIS), Combo 8 | Statewide† | 47.2% 47.5% | 45.8% ▼ At 50th |
| | AMG | 43.3% 48.7% | 45.3% At 50th |
| | CCW | 53.8% 59.9% | 58.9% Above 75th |

| | CHPW | 52.8% | 47.4% | 52.8% | At 75th |
|--|------------|-------|-------|-------|------------------------|
| | MHW | 44.0% | 44.0% | 40.9% | At 50th |
| | UHC | 47.9% | 47.9% | 47.0% | At 75th |
| Childhood Immunization Status (CIS), Combo 9 | Statewide† | 42.1% | 42.5% | 41.9% | At 50th |
| | AMG | 39.4% | 45.3% | 40.9% | At 50th |
| | CCW | 47.2% | 54.3% | 52.3% | Above 75th |
| | CHPW | 48.4% | 43.1% | 48.2% | At 75th |
| | MHW | 38.4% | 38.4% | 37.7% | At 50th |
| | UHC | 44.8% | 44.8% | 44.0% | At 75th |
| Childhood Immunization Status (CIS), Combo 10 | Statewide† | 41.5% | 42.1% | 41.7% | At 50th |
| | AMG | 38.4% | 45.3% | 40.4% | At 50th |
| | CCW | 47.0% | 54.0% | 52.3% | Above 75th |
| | CHPW | 47.7% | 42.8% | 47.9% | At 75th |
| | MHW | 38.0% | 38.0% | 37.5% | Below 50th |
| | UHC | 44.3% | 44.3% | 43.6% | At 75th |
| Immunizations for Adolescents (IMA), Meningococcal | Statewide† | 79.0% | 79.6% | 77.2% | Below 50th |
| | AMG | 70.6% | 73.5% | 74.9% | Below 50th |
| | CCW | 82.7% | 81.8% | 78.4% | Below 50th |
| | CHPW | 83.0% | 81.5% | 81.3% | At 50th |
| | MHW | 79.3% | 80.3% | 76.2% | Below 50th |
| | UHC | 69.6% | 72.5% | 76.9% | Below 50th |
| Immunizations for Adolescents (IMA), Tdap | Statewide† | 86.0% | 86.7% | 86.9% | At 50th |
| | AMG | 85.9% | 82.5% | 85.4% | At 50th |
| | CCW | 88.1% | 91.2% | 86.4% | At 50th |
| | CHPW | 88.3% | 85.9% | 90.0% | At 50th |
| | MHW | 86.9% | 87.3% | 86.6% | At 50th |
| | UHC | 74.2% | 80.0% | 85.4% | At 50th |
| Immunizations for Adolescents (IMA), HPV | Statewide† | 39.0% | 43.7% | 43.2% | Above 50th, Below 75th |
| | AMG | 34.5% | 37.2% | 38.2% | At 50th |
| | CCW | 45.5% | 46.0% | 47.2% | At 75th |
| | CHPW | 48.2% | 44.0% | 46.7% | At 75th |
| | MHW | 35.5% | 44.5% | 43.1% | At 50th |
| | UHC | 32.4% | 38.7% | 36.3% | At 50th |
| Immunizations for Adolescents (IMA), Combo 1 | Statewide† | 76.0% | 77.4% | 75.0% | Below 50th |

| | AMG | 68.4% | 69.3% | 72.5% | Below 50th |
|--|--------------|-------|----------------|-------|------------------------|
| | CCW | 79.3% | 79.3% | 75.7% | Below 50th |
| | CHPW | 79.6% | 78.1% | 79.8% | At 50th |
| | MHW | 76.6% | 78.8% | 74.0% | Below 50th |
| | UHC | 65.2% | 70.1% | 74.0% | Below 50th |
| Immunizations for Adolescents (IMA), Combo 2 | Statewide† | 36.7% | 41.4% | 39.6% | Above 50th, Below 75th |
| | AMG | 32.4% | 33.6% | 36.0% | At 50th |
| | CCW | 42.6% | 41.8% | 42.8% | At 75th |
| | CHPW | 44.8% | 41.1% | 44.0% | At 75th |
| | MHW | 33.8% | 43.3% 🔺 | 38.9% | At 50th |
| | UHC | 30.2% | 35.3% | 34.3% | At 50th |
| Lead Screening in Children (LSC) | Statewide† | 31.7% | 29.8% | 33.7% | Below 50th |
| | AMG | 32.1% | 32.1% | 31.4% | Below 50th |
| | CCW | 36.7% | 37.5% | 33.6% | Below 50th |
| | CHPW | 46.5% | 39.7% | 41.1% | Below 50th |
| | MHW | 27.7% | 25.5% | 34.1% | Below 50th |
| | UHC | 24.1% | 25.5% | 26.0% | Below 50th |
| Breast Cancer Screening (BCS) | Statewide† | 54.5% | 52.0% ▼ | 48.0% | Below 50th |
| | AMG | 47.2% | 45.9% | 42.5% | ▼ Below 50th |
| | CCW | 51.6% | 52.0% | 49.2% | ▼ Below 50th |
| | CHPW | 66.7% | 51.5% ▼ | 46.2% | Below 50th |
| | MHW | 53.9% | 53.8% | 49.7% | Below 50th |
| | UHC | 50.6% | 52.8% | 48.9% | Below 50th |
| Cervical Cancer Screening (CCS) | Statewide† | 57.7% | 60.5% | 58.6% | Below 50th |
| | AMG | 54.1% | 54.1% | 47.7% | Below 50th |
| | CCW | 53.3% | 56.0% | 51.1% | Below 50th |
| | CHPW | 61.6% | 54.0% | 55.0% | At 50th |
| | MHW | 60.8% | 66.2% | 65.9% | At 75th |
| | UHC | 50.5% | 57.7% | 53.3% | Below 50th |
| Chlamydia Screening (CHL), 16-20 Years | Statewide† | 49.2% | 48.9% | 45.5% | |
| | AMG | 46.5% | 47.8% | 45.8% | Below 50th |
| | CCW | 52.0% | 51.0% | 48.2% | Below 50th |
| | CHPW | 48.4% | 46.9% | 45.0% | Below 50th |
| | - | | | | |

| | UHC | 47.6% | 47.4% | 44.0% Below 50th |
|---|------------|-------|-------|------------------------------|
| Chlamydia Screening (CHL), 21-24 Years | Statewide† | 60.1% | 59.5% | 55.2% ▼ Below 50th |
| | AMG | 57.4% | 58.6% | 52.7% ▼ Below 50th |
| | CCW | 60.2% | 61.3% | 57.6% ▼ Below 50th |
| | CHPW | 60.3% | 58.9% | 53.7% ▼ Below 50th |
| | MHW | 61.0% | 60.1% | 56.2% ▼ Below 50th |
| | UHC | 58.3% | 56.3% | 52.2% ▼ Below 50th |
| Chlamydia Screening (CHL), Total | Statewide† | 54.2% | 53.6% | 49.9% ▼ Below 50th |
| | AMG | 52.6% | 53.4% | 49.4% ▼ Below 50th |
| | CCW | 55.7% | 55.4% | 52.2% ▼ Below 50th |
| | CHPW | 53.9% | 52.2% | 48.9% ▼ Below 50th |
| | MHW | 54.4% | 54.0% | 50.2% ▼ Below 50th |
| | UHC | 53.1% | 51.6% | 48.0% ▼ Below 50th |
| Appropriate Testing for Pharyngitis (CWP), 3-17 Years | Statewide† | NR | 79.6% | 80.6% Below 50th |
| | AMG | NR | 76.2% | 75.2% Below 50th |
| | CCW | NR | 75.6% | 78.9% ▲ Below 50th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 79.5% | 80.9% At 50th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 80.8% | 81.2% At 50th |
| | UHC | NR | 81.5% | 84.5% At 75th |
| Appropriate Testing for Pharyngitis (CWP), 18-64 Years | Statewide† | NR | 76.8% | 76.0% Above 75th |
| | AMG | NR | 73.5% | 73.8% At 75th |
| | CCW | NR | 76.3% | 75.3% At 75th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 78.5% | 77.7% Above 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 77.2% | 76.6% Above 75th |
| | UHC | NR | 76.3% | 74.1% At 75th |
| Appropriate Testing for Pharyngitis (CWP), 65+ Years | Statewide† | NR | *** | *** At 50th |
| | AMG | NR | NR | *** At 50th |
| | CCW | NR | *** | *** At 50th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | *** | NR Below 50th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | *** | *** At 50th |
| | UHC | NR | *** | *** At 50th |
| Appropriate Testing for Pharyngitis (CWP), Total | Statewide† | NR | 78.6% | 78.7% Above 50th, Below 75th |
| 7 7 | AMG | NR | 74.9% | 74.5% Below 50th |
| | CCW | NR | 75.9% | 77.6% At 50th |

| Due to substantial changes in measure specifications between MY | CHPW | NR | 79.1% | 79.7% | Above 50th, Below 75th |
|---|------------|-------|---------|----------------|------------------------|
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 79.5% | 79.4% | Above 50th, Below 75th |
| | UHC | NR | 79.2% | 79.3% | Above 50th, Below 75th |
| Use of Spirometry Testing in the Assessment and Diagnosis of | Statewide† | 27.3% | 26.5% | 21.8% 🔻 | Below 50th |
| COPD (SPR) | AMG | 27.7% | 26.3% | 24.1% | At 50th |
| | CCW | 24.5% | 23.5% | 20.7% | Below 50th |
| | CHPW | 29.0% | 28.9% | 21.1% 🔻 | Below 50th |
| | MHW | 28.5% | 26.9% | 22.5% | Below 50th |
| | UHC | 24.6% | 25.1% | 19.3% | Below 50th |
| Pharmacotherapy Management of COPD Exacerbation (PCE), | Statewide† | 74.2% | 74.1% | 71.9% | At 50th |
| Systemic Corticosteroid | AMG | 70.8% | 72.1% | 71.3% | At 50th |
| | CCW | 72.2% | 73.5% | 71.4% | At 50th |
| | CHPW | 76.7% | 74.9% | 74.1% | At 50th |
| | MHW | 75.2% | 74.7% | 73.7% | At 75th |
| | UHC | 73.6% | 74.2% | 68.2% | At 50th |
| Pharmacotherapy Management of COPD Exacerbation (PCE), | Statewide† | 85.8% | 85.1% | 84.8% | At 50th |
| Bronchodilator | AMG | 84.1% | 83.6% | 85.1% | At 50th |
| | CCW | 83.8% | 84.4% | 88.2% | At 50th |
| | CHPW | 87.7% | 83.9% | 87.2% | At 50th |
| | MHW | 86.1% | 86.1% | 86.8% | At 50th |
| | UHC | 85.7% | 86.6% | 78.2% ▼ | Below 50th |
| Asthma Medication Ratio (AMR), 5-11 Years | Statewide† | 66.2% | 65.3% | 74.7% 🔺 | Below 50th |
| | AMG | 65.6% | 59.3% | 81.4% 🔺 | At 50th |
| | CCW | 70.1% | 68.1% | 77.3% | At 50th |
| | CHPW | 59.8% | 55.4% | 60.9% | Below 50th |
| | MHW | 67.1% | 67.3% | 76.5% 🔺 | Below 50th |
| | UHC | 67.7% | 69.2% | 74.7% | At 50th |
| Asthma Medication Ratio (AMR), 12-18 Years | Statewide† | 54.8% | 56.6% | 65.8% 🔺 | Below 50th |
| | AMG | 52.8% | 52.3% | 66.7% | At 50th |
| | CCW | 55.9% | 55.4% | 62.1% | Below 50th |
| | CHPW | 49.7% | 57.3% | 61.2% | Below 50th |
| | MHW | 56.0% | 56.9% | 68.2% 🔺 | At 50th |
| | UHC | 57.1% | 57.3% | 61.8% | Below 50th |
| Asthma Medication Ratio (AMR), 19-50 Years | Statewide† | 45.8% | 49.9% 🔺 | 57.2% 🔺 | At 50th |

| CCW CHPW 49.1% 44.6% 61.6% 61.6% 62.9% Above 75th Above 75th MHW 44.6% 61.6% 61.6% 62.9% Above 75th ABOVE 7 | | | | | |
|---|--|------------|-------|---------|------------------------------|
| CHPW 44.6% 61.6% | | AMG | 44.0% | 44.0% | 51.7% 🔺 Below 50th |
| MHW 46.3% 47.6% 57.5% ♣ At 50th UHC 44.3% 47.1% 52.9% Below 50th Asthma Medication Ratio (AMR), 51-64 Years Statewide† 51.3% 54.5% 59.7% ♣ At 75th AMG 54.1% 50.8% 56.0% At 50th CCW 53.1% 55.5% 57.7% At 50th CHPW 52.4% 63.0% ♣ 64.8% At 75th AMHW 50.3% 51.7% 60.2% ♣ At 75th Asthma Medication Ratio (AMR), Total Statewide† 52.7% 55.0% ♣ 64.8% At 50th Asthma Medication Ratio (AMR), Total Statewide† 52.7% 55.0% ♣ 62.1% ♠ Below 50th AMG 49.6% 48.2% 57.8% ♠ Below 50th CCW 55.6% 56.0% ♠ 60.7% ♠ Below 50th AMG 49.6% 48.2% 57.8% ♠ Below 50th CHPW 50.1% 59.9% ♠ 64.2% ♠ At 50th Controlling High Blood Pressure (CBP) Statewide† NR NR 57.4% At 50th COW NR NR S7.4% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR S8.4% At 50th Due to substantial changes in measure results are MHW NR NR 63.3% At 75th Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 99.6% 82.2% 83.9% At 50th AMG 67.3% 80.4% 77.0% At 50th AMG 67.3% 83.8% 84.0% At 75th AMG 67.3% 83.8% 84.0% At 75th AMG 83.8% 83.8% 84.0% At 75th AND COW 79.8% 83.8% 84.0% At 75th AMG 83.8% 83.5% 83.8% 84.50th AMG 83.8% 83.5% 83.8% 84.50th AMG 83.8% 83.5% 83.8% 84.50th At 50th AMG 83.8% 83.5% 83.8% 84.50th At 50th AMG 83.8% 83.5% 83.8% 84.50th At 50th At 50th | | CCW | 49.1% | 49.9% | 53.7% At 50th |
| Here | | CHPW | 44.6% | 61.6% | 66.2% Above 75th |
| Asthma Medication Ratio (AMR), 51-64 Years AMG CCW 53.1% CHPW 52.4% AMG At 75th At 50th CHPW 52.4% At 50th Asthma Medication Ratio (AMR), Total At 50th AMG AP9.6% AB.2% 57.8% Below 50th AMG CCW 55.6% 56.0% 60.7% AB Below 50th CHPW 50.1% 59.9% A 64.2% At 50th At 50th At 50th CONTrolling High Blood Pressure (CBP) Statewide† AMG NR NR S8.6% Above 50th, Below 50th AMG CCW NR NR S8.4% At 50th | | MHW | 46.3% | 47.6% | 57.5% 🔺 At 50th |
| AMG 54.1% 50.8% 56.0% At 50th CCW 53.1% 55.5% 57.7% At 50th CHPW 52.4% 63.0% | | UHC | 44.3% | 47.1% | 52.9% Below 50th |
| CCW 53.1% 55.5% 57.7% At 50th CHPW 52.4% 63.0% ▲ 64.8% At 75th MHW 50.3% 51.7% 60.2% ▲ At 75th UHC 49.4% 53.7% 57.4% At 50th Asthma Medication Ratio (AMR), Total Statewide† 52.7% 55.0% ▲ 62.1% ▲ Below 50th AMG 49.6% 48.2% 57.8% ▲ Below 50th CCW 55.6% 56.0% 60.7% ▲ Below 50th CHPW 50.1% 59.9% ▲ 64.2% ▲ At 50th MHW 53.9% 54.6% 63.7% ▲ Below 50th UHC 50.6% 53.3% 57.7% Below 50th Controlling High Blood Pressure (CBP) Statewide† NR NR 58.6% Above 50th, Below 50th CCW NR NR S8.6% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR 58.4% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR 63.3% At 75th 2019 and MY 2020, only current year measure results are displayed. UHC NR NR 62.3% At 75th Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CCW 79.8% 83.8% 78.3% At 50th MHW 79.3% 79.5% 86.9% At 50th Statewide† 77.5% 79.6% 82.2% 83.9% At 50th CCW 79.8% 83.8% 78.3% At 50th MHW 79.3% 79.5% 86.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 83.8% 84.0% At 75th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.8% 83.8% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | Asthma Medication Ratio (AMR), 51-64 Years | Statewide† | 51.3% | 54.5% | 59.7% 🔺 At 75th |
| CHPW MHW S0.3% S1.7% 60.2% A At 75th C15th | | AMG | 54.1% | 50.8% | 56.0% At 50th |
| MHW 50.3% 51.7% 60.2% ▲ At 75th UHC 49.4% 53.7% 57.4% At 50th Asthma Medication Ratio (AMR), Total Statewide† 52.7% 55.0% ▲ 62.1% ▲ Below 50th AMG 49.6% 48.2% 57.8% ▲ Below 50th CWW 55.6% 56.0% 60.7% ▲ Below 50th CWW 50.1% 59.9% ▲ 64.2% ▲ At 50th CWW 50.6% 53.3% 57.7% Below 50th CWW 50.6% 53.3% 57.7% At 50th CWW 50.6% 53.3% 57.7% 58.4% 50th CWW 50.6% 53.3% 57.7% 58.4% 50th CWW 50.6% 53.3% 57.7% 50th CWW 50.6% 50.6% 50th CWW 50.6% 50t | | CCW | 53.1% | 55.5% | 57.7% At 50th |
| UHC 49.4% 53.7% 57.4% At 50th Asthma Medication Ratio (AMR), Total Statewide† 52.7% 55.0% ▲ 62.1% ▲ Below 50th AMG 49.6% 48.2% 57.8% ▲ Below 50th CCW 55.6% 56.0% 60.7% ▲ Below 50th CHPW 50.1% 59.9% ▲ 64.2% ▲ At 50th MHW 53.9% 54.6% 63.7% ▲ Below 50th Controlling High Blood Pressure (CBP) Statewide† NR NR 57.7% Below 50th Controlling High Blood Pressure (CBP) Statewide† NR NR 58.6% Above 50th, Below 75t AMG NR NR NR 57.4% At 50th CCW NR NR NR 58.6% Above 50th, Below 75t AMG NR NR NR 58.6% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR NR 58.4% At 75th Persistence of Beta-Blocker Treatment After a | | CHPW | 52.4% | 63.0% 🔺 | 64.8% At 75th |
| Asthma Medication Ratio (AMR), Total AMG AMG AMG AMG AMG AMG AMG AM | | MHW | 50.3% | 51.7% | 60.2% 🔺 At 75th |
| AMG 49.6% 48.2% 57.8% ♣ Below 50th CCW 55.6% 56.0% 60.7% ♣ Below 50th CHPW 50.1% 59.9% ♣ 64.2% ♠ At 50th MHW 53.9% 54.6% 63.7% ♣ Below 50th MHW 53.9% 54.6% 63.7% ♣ Below 50th MHW 53.9% 54.6% 63.7% ♣ Below 50th DHC 50.6% 53.3% 57.7% Below 50th Controlling High Blood Pressure (CBP) Statewide† NR NR NR 58.6% Above 50th, Below 75th AMG NR NR S8.6% Above 50th, Below 75th CCW NR NR NR 58.4% At 50th CCW NR NR S8.4% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR 63.3% At 75th 2019 and MY 2020, only current year measure results are MHW NR NR 56.0% At 50th Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 79.6% 82.7% At 50th AMG 67.3% 80.4% 77.0% At 50th AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th OHPC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 83.8% At 50th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | | UHC | 49.4% | 53.7% | 57.4% At 50th |
| CCW 55.6% 56.0% 60.7% ♣ Below 50th CHPW 50.1% 59.9% ♣ 64.2% ♠ At 50th MHW 53.9% 54.6% 63.7% ♣ Below 50th MHW 53.9% 54.6% 63.7% ♣ Below 50th UHC 50.6% 53.3% 57.7% Below 50th Controlling High Blood Pressure (CBP) Statewide† NR NR S8.6% Above 50th, Below 75th AMG NR NR S7.4% At 50th CCW NR NR NR 58.4% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR S8.4% At 50th 2019 and MY 2020, only current year measure results are MHW NR NR S6.0% At 75th displayed. UHC NR NR S6.0% At 50th Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 79.6% 82.7% At 50th AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 83.5% 83.8% At 50th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | Asthma Medication Ratio (AMR), Total | Statewide† | 52.7% | 55.0% 🔺 | 62.1% ▲ Below 50th |
| CHPW 50.1% 59.9% ▲ 64.2% ▲ At 50th MHW 53.9% 54.6% 63.7% ▲ Below 50th UHC 50.6% 53.3% 57.7% Below 50th Controlling High Blood Pressure (CBP) Statewide† NR NR NR 58.6% Above 50th, Below 75t AMG NR NR NR 57.4% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR 58.6% At 50th 2019 and MY 2020, only current year measure results are displayed. MHW NR NR NR 56.0% At 50th Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 79.6% 82.7% At 50th AMG 67.3% 80.4% 77.0% At 50th CW 79.8% 83.8% 78.3% At 50th MHW 79.6% 82.2% 83.9% At 50th MHW 79.6% 82.2% 83.9% At 50th WHW 79.6% 82.2% 83.9% At 50th WHW 79.6% < | | AMG | 49.6% | 48.2% | 57.8% A Below 50th |
| MHW 53.9% 54.6% 63.7% ▲ Below 50th UHC 50.6% 53.3% 57.7% Below 50th UHC 50.6% 53.3% 57.7% Below 50th Controlling High Blood Pressure (CBP) Statewide† NR NR S8.6% Above 50th, Below 75th AMG NR NR 57.4% At 50th CCW NR NR S8.6% At 50th CCW NR NR S8.6% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR 63.3% At 75th 2019 and MY 2020, only current year measure results are MHW NR NR 56.0% At 50th displayed. UHC NR NR 62.3% At 75th Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 79.6% 82.7% At 50th AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 84.0% At 75th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th | | CCW | 55.6% | 56.0% | 60.7% ▲ Below 50th |
| UHC 50.6% 53.3% 57.7% Below 50th | | CHPW | 50.1% | 59.9% 🔺 | 64.2% 🔺 At 50th |
| Controlling High Blood Pressure (CBP) Statewide† NR NR NR S8.6% Above 50th, Below 75th AMG NR NR NR NR S7.4% At 50th CCW NR NR NR S8.4% At 50th NR Due to substantial changes in measure specifications between MY CHPW NR NR NR NR S6.0% At 75th At 50th NR NR R S6.0% At 50th NR NR NR S6.0% At 50th NR NR NR S6.0% At 50th NR NR NR S6.0% At 50th AMG G7.3% S0.4% 77.0% At 50th CCW 79.8% S3.8% 78.3% At 50th CHPW 79.6% S2.2% S3.9% At 50th MHW 79.3% 79.5% S6.9% At 50th NHW 79.3% 79.5% S6.9% At 50th NHW 79.3% 79.5% S6.9% At 50th NHW 79.3% 79.5% S6.9% At 75th NHW NR NR S6.0% At 50th NR NR S6.0% At 50th NR NR NR NR NR NR S6.0% At 50th NR NR NR NR NR NR S6.0% At 50th NR NR NR NR NR NR S6.0% At 50th NR NR NR NR NR NR S6.0% At 50th NR NR NR NR NR NR NR NR S6.0% At 50th NR NR NR NR NR NR NR S6.0% At 50th NR NR NR NR NR NR NR S6.0% At 50th NR NR NR NR NR NR NR S6.0% At 50th NR NR NR NR NR NR NR S6.0% At 75th NR NR NR NR NR NR NR NR NR N | | MHW | 53.9% | 54.6% | 63.7% 🔺 Below 50th |
| AMG NR NR 57.4% At 50th CCW NR NR NR 58.4% At 50th Due to substantial changes in measure specifications between MY CHPW NR NR NR 63.3% At 75th 2019 and MY 2020, only current year measure results are MHW NR NR NR 56.0% At 50th displayed. UHC NR NR NR 62.3% At 75th Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 79.6% 82.7% At 50th AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th CT 75.5% TO 10 The approximate of the substantial changes in measure specifications between MY CHPW NR NR 63.3% At 75th CCHPW 79.6% 82.2% 83.9% At 50th CCHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th CT 75.5% TO 10 The approximate of the substantial changes in measure specifications between MY CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th TO 10 The approximate of the substantial changes in measure specifications between MY CHPW 79.6% 82.2% 83.8% At 50th MHW 79.3% 79.5% 86.9% At 75th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 83.8% 84.0% At 50th CCW 82.3% 80.0% 81.7% At 50th | | UHC | 50.6% | 53.3% | 57.7% Below 50th |
| CCW | Controlling High Blood Pressure (CBP) | Statewide† | NR | NR | 58.6% Above 50th, Below 75th |
| Due to substantial changes in measure specifications between MY CHPW NR NR S6.33% At 75th 2019 and MY 2020, only current year measure results are MHW NR NR S6.0% At 50th displayed. Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 79.6% 82.7% At 50th AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 83.8% 84.0% At 75th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | | AMG | NR | NR | 57.4% At 50th |
| 2019 and MY 2020, only current year measure results are MHW NR NR 56.0% At 50th displayed. Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th CCW | | CCW | NR | NR | 58.4% At 50th |
| displayed. UHC NR NR 62.3% At 75th Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 79.6% 82.7% At 50th AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 84.0% At 75th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th | Due to substantial changes in measure specifications between MY | CHPW | NR | NR | 63.3% At 75th |
| Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) Statewide† 77.5% 79.6% 82.7% At 50th AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 84.0% At 75th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | 2019 and MY 2020, only current year measure results are | MHW | NR | NR | 56.0% At 50th |
| AMG 67.3% 80.4% 77.0% At 50th CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | displayed. | UHC | NR | NR | 62.3% At 75th |
| CCW 79.8% 83.8% 78.3% At 50th CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) | Statewide† | 77.5% | 79.6% | 82.7% At 50th |
| CHPW 79.6% 82.2% 83.9% At 50th MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | | AMG | 67.3% | 80.4% | 77.0% At 50th |
| MHW 79.3% 79.5% 86.9% At 75th UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 84.0% At 75th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | | CCW | 79.8% | 83.8% | 78.3% At 50th |
| UHC 76.4% 72.3% 77.9% At 50th Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 84.0% At 75th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | | CHPW | 79.6% | 82.2% | 83.9% At 50th |
| Statin Therapy for Patients With Cardiovascular Disease (SPC), Statewide† 83.3% 83.8% 84.0% At 75th Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | | MHW | 79.3% | 79.5% | 86.9% At 75th |
| Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | | UHC | 76.4% | 72.3% | 77.9% At 50th |
| Received Statin Therapy, 21-75 years (Male) AMG 83.8% 83.5% 83.8% At 50th CCW 82.3% 80.0% 81.7% At 50th | Statin Therapy for Patients With Cardiovascular Disease (SPC), | | 83.3% | 83.8% | |
| CCW 82.3% 80.0% 81.7% At 50th | Received Statin Therapy, 21-75 years (Male) | AMG | 83.8% | 83.5% | 83.8% At 50th |
| | | CCW | 82.3% | 80.0% | 81.7% At 50th |
| | | CHPW | 81.0% | 83.7% | 80.9% At 50th |
| MHW 85.6% 84.8% 85.6% At 75th | | MHW | | | |

| | UHC | 81.6% | 84.9% | 84.2% At 50th |
|--|------------|-------|-------|--------------------------------|
| Statin Therapy for Patients With Cardiovascular Disease (SPC), | Statewide† | 82.0% | 82.0% | 81.3% At 75th |
| Received Statin Therapy, 40-75 years (Female) | AMG | 82.5% | 78.1% | 79.8% At 50th |
| | CCW | 83.2% | 85.1% | 81.3% At 50th |
| | CHPW | 82.1% | 83.3% | 85.6% At 75th |
| | MHW | 81.5% | 80.8% | 80.5% At 50th |
| | UHC | 82.1% | 84.1% | 81.4% At 50th |
| Statin Therapy for Patients With Cardiovascular Disease (SPC), | Statewide† | 82.8% | 83.2% | 83.1% At 75th |
| Received Statin Therapy Total | AMG | 83.4% | 82.1% | 82.6% At 50th |
| | CCW | 82.6% | 81.8% | 81.6% At 50th |
| | CHPW | 81.4% | 83.6% | 82.3% At 50th |
| | MHW | 84.0% | 83.3% | 83.8% At 75th |
| | UHC | 81.8% | 84.7% | 83.2% At 75th |
| Statin Therapy for Patients With Cardiovascular Disease (SPC), | Statewide† | 65.7% | 68.3% | 74.3% 🛕 Above 50th, Below 75th |
| Statin Adherence 80%, 21-75 years (Male) | AMG | 64.8% | 70.7% | 70.6% At 50th |
| | CCW | 67.6% | 66.7% | 71.5% At 50th |
| | CHPW | 68.3% | 71.6% | 80.4% 🛕 At 75th |
| | MHW | 65.7% | 68.5% | 76.1% 🔺 At 75th |
| | UHC | 62.5% | 63.0% | 70.3% At 50th |
| Statin Therapy for Patients With Cardiovascular Disease (SPC), | Statewide† | 68.2% | 70.8% | 75.9% 🛕 At 75th |
| Statin Adherence 80%, 40-75 years (Female) | AMG | 66.2% | 65.9% | 69.9% At 50th |
| | CCW | 71.3% | 73.3% | 70.4% At 50th |
| | CHPW | 66.5% | 74.0% | 83.1% At 75th |
| | MHW | 67.9% | 70.3% | 76.8% At 75th |
| | UHC | 69.7% | 69.0% | 76.0% At 50th |
| Statin Therapy for Patients With Cardiovascular Disease (SPC), | Statewide† | 66.6% | 69.1% | 74.9% 🛕 Above 50th, Below 75th |
| Statin Adherence 80%, Total | AMG | 65.2% | 69.5% | 70.4% At 50th |
| | CCW | 68.9% | 69.1% | 71.1% At 50th |
| | CHPW | 67.6% | 72.5% | 81.3% 🛕 Above 75th |
| | MHW | 66.5% | 69.2% | 76.3% 🛕 At 75th |
| | UHC | 64.8% | 64.8% | 72.3% 🛕 At 50th |
| Comprehensive Diabetes Care (CDC), HbA1c Testing | Statewide† | 89.5% | 89.5% | 84.7% ▼ At 75th |
| | AMG | 86.6% | 90.3% | 87.8% At 75th |
| | CCW | 88.6% | 90.0% | 85.4% At 50th |

| | CHDW | 00.00/ | 90.20/ | 01 10/ | ۸+ ۵+ ۱ |
|---|-------------|--------|----------------|----------------|------------------------|
| | CHPW MHW | 90.0% | 89.3% 88.8% | 84.4% 83.0% | At 50th At 50th |
| | | 90.0% | | | |
| Community District Community (CDC) 2 | UHC | 90.5% | 90.3% | 86.4% | At 50th |
| Comprehensive Diabetes Care (CDC), Poor HbA1c Control (Note | Statewide† | 37.1% | 34.5% | 37.5% | Above 75th |
| that a lower score is better for this measure) | AMG | 39.7% | 36.5% | 40.2% | Above 75th |
| | CCW | 47.0% | 41.4% | 45.3% | Above 50th, Below 75th |
| | CHPW | 43.1% | 38.4% | 41.1% | Above 75th |
| | MHW | 32.6% | 31.6% | 36.3% | Above 75th |
| | UHC | 31.9% | 30.9% | 30.4% | Above 75th |
| Comprehensive Diabetes Care (CDC), HbA1c Control < 8.0% | Statewide† | 50.3% | 51.9% | 51.9% | At 75th |
| | AMG | 47.2% | 51.8% | 47.7% | At 50th |
| | CCW | 42.1% | 47.0% | 44.5% | At 50th |
| | CHPW | 44.3% | 46.7% | 48.9% | At 50th |
| | MHW | 54.0% | 54.0% | 53.5% | At 75th |
| | UHC | 56.4% | 56.2% | 58.2% | Above 75th |
| Comprehensive Diabetes Care (CDC), Eye Exam | Statewide† | 58.5% | 59.1% | 51.6% | At 50th |
| | AMG | 51.8% | 50.1% | 49.6% | At 50th |
| | CCW | 56.4% | 61.1% | 51.6% | At 50th |
| | CHPW | 57.9% | 58.2% | 54.0% | At 50th |
| | MHW | 60.6% | 61.6% | 50.6% | At 50th |
| | UHC | 60.1% | 59.1% | 54.0% | At 50th |
| Comprehensive Diabetes Care (CDC), Blood Pressure Control < | Statewide† | NR | NR | 68.4% | At 75th |
| 140/90 mm Hg | AMG | NR | NR | 64.7% | At 75th |
| | CCW | NR | NR | 65.7% | At 75th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | NR | 66.2% | At 75th |
| 2019 and MY 2020, only current year measure results are | MHW | NR | NR | 70.1% | At 75th |
| displayed. | UHC | NR | NR | 70.6% | Above 75th |
| Statin Therapy for Patients With Diabetes (SPD), Received Statin | Statewide† | 64.4% | 66.0% 🔺 | 66.4% | At 50th |
| Therapy | AMG | 63.5% | 65.2% | 65.8% | At 50th |
| • • | CCW | 64.0% | 64.8% | 65.9% | At 50th |
| | CHPW | 65.3% | 67.9% | 66.1% | At 50th |
| | MHW | 64.2% | 65.6% | 66.0% | At 50th |
| | UHC | 64.9% | 66.4% | 68.6% | At 75th |
| Statin Therapy for Patients With Diabetes (SPD), Statin Adherence | | 63.9% | 66.7% 🛕 | 73.1% 🔺 | |

| 80% | AMG | | 61.4% | 66 | .8% | | 68.29 | 6 | At 50th |
|--|--------------|-----|-------|----|------|----------|-------|------------|------------------------|
| | CCW | | 65.4% | 69 | .5% | | 72.09 | 6 | At 75th |
| | CHPW | | 67.4% | 71 | .1% | | 76.69 | 6 | Above 75th |
| | MHW | | 63.1% | 64 | .4% | | 74.59 | 6 | Above 75th |
| | UHC | | 62.2% | 64 | .8% | | 71.29 | 6 | Above 50th, Below 75th |
| Antidepressant Medication Management (AMM), Effective Acute | F Statewide† | | 50.9% | 53 | 3.5% | A | 58.59 | 6 ▲ | Above 50th, Below 75th |
| | AMG | | 51.0% | 52 | .9% | | 52.99 | 6 | Below 50th |
| | CCW | | 54.8% | 56 | 5.2% | | 58.99 | 6 | Above 50th, Below 75th |
| | CHPW | | 51.0% | 55 | .3% | A | 57.49 | 6 | At 50th |
| | MHW | | 49.2% | 52 | .4% | A | 60.69 | 6 | Above 50th, Below 75th |
| | UHC | | 52.2% | 53 | .6% | | 57.19 | 6 | At 50th |
| Antidepressant Medication Management (AMM), Continuation | Statewide† | | 36.0% | 38 | 3.4% | A | 42.99 | 6 ▲ | Above 50th, Below 75th |
| Phase | AMG | | 36.1% | 39 | .7% | A | 38.49 | 6 | Below 50th |
| | CCW | | 38.1% | 39 | .9% | | 41.19 | 6 | At 50th |
| | CHPW | | 36.8% | 39 | .9% | A | 42.69 | 6 | Above 50th, Below 75th |
| | MHW | | 34.4% | 37 | .2% | A | 44.69 | 6 ▲ | Above 50th, Below 75th |
| | UHC | | 38.0% | 38 | 3.8% | | 42.69 | 6 | Above 50th, Below 75th |
| Follow-Up Care for Children Prescribed ADHD Medication (ADD), | Statewide† | | 42.8% | 43 | .9% | | 45.29 | 6 | Below 50th |
| Initiation | AMG | | 36.5% | 34 | .0% | | 44.59 | 6 ▲ | Below 50th |
| | CCW | | 38.6% | 39 | .6% | | 46.29 | 6 | Below 50th |
| | CHPW | | 40.1% | 46 | .2% | | 41.59 | 6 | Below 50th |
| | MHW | | 46.4% | 47 | '.1% | | 45.49 | 6 | Below 50th |
| | UHC | | 37.2% | 37 | '.4% | | 47.39 | 6 ▲ | Below 50th |
| Follow-Up Care for Children Prescribed ADHD Medication (ADD), | Statewide† | | 50.8% | 53 | .6% | | 52.49 | 6 | Above 75th |
| Continuation | AMG | | 45.5% | 47 | '.3% | | 52.89 | 6 | At 50th |
| | CCW | | 44.1% | 48 | 3.9% | | 53.69 | 6 | At 75th |
| | CHPW | | 48.2% | 56 | .7% | | 44.79 | 6 | At 50th |
| | MHW | | 55.7% | 58 | 3.3% | | 53.09 | 6 | Above 75th |
| | UHC | | 46.4% | 37 | '.5% | | 53.79 | 6 | At 75th |
| Follow-Up after Hospitalization for Mental Illness (FUH), 30-Day | Statewide† | | 88.2% | 63 | 3.3% | ▼ | 74.29 | 6 | Above 50th, Below 75th |
| Follow-Up, 6-17 Years | AMG | NR | | 61 | 7% | | 74.19 | 6 | At 50th |
| | CCW | *** | | 52 | .2% | | 68.99 | 6 | At 50th |
| | CHPW | *** | | 54 | .2% | | 69.89 | 6 ▲ | At 50th |
| | MHW | | 89.4% | 79 | .0% | | 79.89 | 6 | At 75th |
| | | | | | | | | | |

| | UHC | NR | | 31.3% | 65.3% 🔺 | At 50th |
|--|------------|-----|-------|----------------|---------|------------------------|
| Follow-Up after Hospitalization for Mental Illness (FUH), 30-Day | Statewide† | | 48.5% | 44.6% | 53.5% 🔺 | At 50th |
| Follow-Up, 18-64 Years | AMG | *** | | 34.2% ▼ | 42.8% 🔺 | Below 50th |
| | CCW | *** | | 34.2% ▼ | 36.0% | Below 50th |
| | CHPW | | 46.2% | 40.6% | 60.0% 🔺 | Above 50th, Below 75th |
| | MHW | | 47.8% | 58.6% ^ | 63.5% 🔺 | At 75th |
| | UHC | NR | | 26.1% | 44.6% 🔺 | Below 50th |
| Follow-Up after Hospitalization for Mental Illness (FUH), 30-Day | Statewide† | NR | | *** | *** | Below 50th |
| Follow-Up, 65+ Years | AMG | NR | | NR | NR | Below 50th |
| | CCW | NR | | NR | *** | At 50th |
| | CHPW | NR | | *** | *** | At 50th |
| | MHW | NR | | NR | *** | At 50th |
| | UHC | NR | | NR | *** | At 50th |
| Follow-Up after Hospitalization for Mental Illness (FUH), 30-Day | Statewide† | | 52.1% | 48.3% | 57.2% 🔺 | Below 50th |
| Follow-Up, Total | AMG | *** | | 37.3% | 46.6% 🔺 | Below 50th |
| | CCW | *** | | 40.7% ▼ | 47.0% 🔺 | Below 50th |
| | CHPW | | 49.1% | 42.5% | 61.3% 🔺 | At 50th |
| | MHW | | 52.0% | 62.8% 🔺 | 66.8% 🔺 | At 75th |
| | UHC | NR | | 26.8% | 46.9% 🔺 | Below 50th |
| Follow-Up after Hospitalization for Mental Illness (FUH), 7-Day | Statewide† | | 69.7% | 44.8% ▼ | 55.7% 🔺 | Above 50th, Below 75th |
| Follow-Up, 6-17 Years | AMG | NR | | 48.6% | 57.5% | At 75th |
| | CCW | *** | | 35.7% | 51.9% 🔺 | At 50th |
| | CHPW | *** | | 35.2% | 47.8% | At 50th |
| | MHW | | 71.2% | 56.9% | 59.9% | At 75th |
| | UHC | NR | | 20.5% | 51.5% 🔺 | At 50th |
| Follow-Up after Hospitalization for Mental Illness (FUH), 7-Day | Statewide† | | 31.5% | 28.9% | 36.7% 🔺 | Above 50th, Below 75th |
| Follow-Up, 18-64 Years | AMG | *** | | 21.3% 🔻 | 27.6% 🔺 | Below 50th |
| | CCW | *** | | 22.8% 🔻 | 25.2% | Below 50th |
| | CHPW | | 30.8% | 24.4% | 40.3% 🔺 | Above 50th, Below 75th |
| | MHW | | 30.4% | 38.0% 🔺 | 44.5% 🔺 | At 75th |
| | UHC | NR | | 19.1% | 30.7% 🔺 | Below 50th |
| Follow-Up after Hospitalization for Mental Illness (FUH), 7-Day | Statewide† | NR | | *** | *** | At 50th |
| Follow-Up, 65+ Years | AMG | NR | | NR | NR | Below 50th |
| | CCW | NR | | NR | *** | At 50th |

| | CHPW | NR | | *** | *** | At 50th |
|---|------------|-----|-------|----------------|---------|------------------------|
| | MHW | NR | | NR | *** | At 50th |
| | UHC | NR | | NR | *** | At 50th |
| Follow-Up after Hospitalization for Mental Illness (FUH), 7-Day | Statewide† | | 35.1% | 32.0% | 40.2% 🔺 | Above 50th, Below 75th |
| Follow-Up, Total | AMG | *** | | 24.3% | 31.1% | Below 50th |
| | CCW | *** | | 27.5% | 34.2% 🔺 | Below 50th |
| | CHPW | | 33.9% | 26.0% | 41.3% | At 50th |
| | MHW | | 34.5% | 41.9% | 47.6% | At 75th |
| | UHC | NR | | 19.3% | 33.1% 🔺 | Below 50th |
| Follow-Up After Emergency Department Visit for Mental Illness | Statewide† | | 56.4% | 66.0% 🔺 | 73.3% 🔺 | Above 50th, Below 75th |
| (FUM), 30-Day Follow-Up, 6-17 Years | AMG | | 33.7% | 49.8% 🔺 | 60.4% | Below 50th |
| | CCW | *** | | 65.3% | 72.6% | At 50th |
| | CHPW | *** | | 63.5% | 62.7% | Below 50th |
| | MHW | | 68.6% | 71.1% | 78.5% 🔺 | At 75th |
| | UHC | NR | | 63.9% | 72.1% | At 50th |
| Follow-Up After Emergency Department Visit for Mental Illness | Statewide† | | 29.3% | 47.3% 🔺 | 53.7% 🔺 | Above 50th, Below 75th |
| (FUM), 30-Day Follow-Up, 18-64 Years | AMG | | 20.9% | 37.9% 🔺 | 42.8% | Below 50th |
| | CCW | | 57.7% | 40.7% ▼ | 38.1% | Below 50th |
| | CHPW | | 56.4% | 57.1% | 61.1% | At 75th |
| | MHW | | 32.2% | 49.2% 🔺 | 59.5% 🔺 | At 75th |
| | UHC | | NR | 49.5% | 53.7% | Above 50th, Below 75th |
| Follow-Up After Emergency Department Visit for Mental Illness | Statewide† | | *** | *** | *** | At 50th |
| (FUM), 30-Day Follow-Up, 65+ Years | AMG | | *** | *** | *** | At 50th |
| | CCW | | NR | NR | *** | At 50th |
| | CHPW | | NR | *** | *** | At 50th |
| | MHW | | NR | NR | NR | Below 50th |
| | UHC | | NR | *** | *** | At 50th |
| Follow-Up After Emergency Department Visit for Mental Illness | Statewide† | | 31.9% | 51.0% 🔺 | 57.8% 🔺 | Above 50th, Below 75th |
| (FUM), 30-Day Follow-Up, Total | AMG | | 22.1% | 39.4% | 45.2% | Below 50th |
| | CCW | | 63.5% | 49.8% | 49.9% | Below 50th |
| | CHPW | | 55.8% | 58.0% | 61.3% | Above 50th, Below 75th |
| | MHW | | 35.5% | 53.6% | 64.0% | At 75th |
| | UHC | | NR | 51.1% | 56.5% | Above 50th, Below 75th |
| Follow-Up After Emergency Department Visit for Mental Illness | Statewide† | | 42.6% | 50.9% 🔺 | 58.2% 🔺 | Above 50th, Below 75th |

| (FUM), 7-Day Follow-Up, 6-17 Years | AMG | 19.2% | 33.2% 🔺 | 44.2% At 50th |
|---|------------|-------|---------|--------------------------------|
| , , , , , , , , , , , , , , , , , , , | CCW | *** | 47.4% | 55.5% Above 50th, Below 75th |
| | CHPW | *** | 52.9% | 48.1% At 50th |
| | MHW | 55.2% | 57.4% | 63.9% 🔺 At 75th |
| | UHC | NR | 45.4% | 59.6% At 75th |
| Follow-Up After Emergency Department Visit for Mental Illness | Statewide† | 17.1% | 34.3% 🔺 | 41.6% Above 50th, Below 75th |
| (FUM), 7-Day Follow-Up, 18-64 Years | AMG | 11.5% | 26.8% 🔺 | 30.7% Below 50th |
| | CCW | 48.1% | 28.0% ▼ | 27.8% Below 50th |
| | CHPW | 34.3% | 42.8% | 48.5% ▲ Above 75th |
| | MHW | 19.0% | 35.7% | 47.5% ▲ Above 75th |
| | UHC | NR | 36.4% | 40.3% Above 50th, Below 75th |
| Follow-Up After Emergency Department Visit for Mental Illness | Statewide† | *** | *** | *** At 50th |
| (FUM), 7-Day Follow-Up, 65+ Years | AMG | *** | *** | *** At 50th |
| | CCW | NR | NR | *** At 50th |
| | CHPW | NR | *** | *** At 50th |
| | MHW | NR | NR | NR Below 50th |
| | UHC | NR | *** | *** At 50th |
| Follow-Up After Emergency Department Visit for Mental Illness | Statewide† | 19.6% | 37.5% 🔺 | 45.1% Above 50th, Below 75th |
| (FUM), 7-Day Follow-Up, Total | AMG | 12.3% | 27.6% | 32.5% ▲ Below 50th |
| | CCW | 52.7% | 35.3% ▼ | 37.2% At 50th |
| | CHPW | 35.0% | 44.1% | 48.4% At 75th |
| | MHW | 22.2% | 40.1% 🔺 | 51.4% 🔺 Above 75th |
| | UHC | NR | 37.4% | 43.2% ▲ Above 50th, Below 75th |
| Follow-Up After Emergency Department Visit for Alcohol and | Statewide† | 20.0% | 14.6% | 17.0% At 75th |
| Other Drug Abuse Dependencies (FUA), 30-Day Follow-Up, 13-17 | AMG | *** | 15.4% | 23.5% At 75th |
| Years | CCW | *** | 14.9% | 15.6% At 75th |
| | CHPW | *** | 17.8% | 20.6% At 75th |
| | MHW | 20.5% | 13.1% | 16.0% At 75th |
| | UHC | NR | *** | 14.8% At 50th |
| Follow-Up After Emergency Department Visit for Alcohol and | Statewide† | 14.9% | 26.5% 🔺 | 29.2% 🔺 Above 75th |
| Other Drug Abuse Dependencies (FUA), 30-Day Follow-Up, 18+ | AMG | 20.0% | 23.5% | 26.7% At 75th |
| Years | CCW | 32.3% | 21.4% | 21.8% At 50th |
| | CHPW | 29.6% | 31.8% | 34.4% Above 75th |
| | MHW | 13.8% | 28.4% 🔺 | 31.6% ▲ Above 75th |

| | UHC | NR | 21.0% | 25.9% 🔺 At 75th |
|---|------------|-------|---------|------------------------------|
| ollow-Up After Emergency Department Visit for Alcohol and | Statewide† | 15.0% | 26.1% 🔺 | 28.7% Above 75th |
| Other Drug Abuse Dependencies (FUA), 30-Day Follow-Up, Tota | l AMG | 19.5% | 23.3% | 26.7% At 75th |
| | CCW | 30.8% | 20.8% | 21.3% At 50th |
| | CHPW | 29.5% | 31.3% | 33.9% Above 75th |
| | MHW | 13.9% | 27.8% | 30.9% ▲ Above 75th |
| | UHC | NR | 21.0% | 25.6% 🔺 At 75th |
| ollow-Up After Emergency Department Visit for Alcohol and | Statewide† | 12.9% | 9.1% | 11.0% At 50th |
| Other Drug Abuse Dependencies (FUA), 7-Day Follow-Up, 13-17 | AMG | *** | 12.8% | 17.7% At 50th |
| ears | CCW | *** | 6.7% | 10.2% At 50th |
| | CHPW | *** | 12.3% | 12.7% At 50th |
| | MHW | 13.7% | 8.6% | 10.3% At 50th |
| | UHC | NR | *** | 8.2% At 50th |
| ollow-Up After Emergency Department Visit for Alcohol and | Statewide† | 8.5% | 16.9% 🔺 | 19.1% 🔺 Above 75th |
| Other Drug Abuse Dependencies (FUA), 7-Day Follow-Up, 18+ | AMG | 12.5% | 15.5% | 18.1% At 75th |
| ears | CCW | 16.1% | 12.6% | 13.6% At 50th |
| | CHPW | 18.9% | 21.2% | 22.0% Above 75th |
| | MHW | 7.8% | 18.1% 🔺 | 20.9% ▲ Above 75th |
| | UHC | NR | 12.2% | 16.6% 🔺 At 75th |
| ollow-Up After Emergency Department Visit for Alcohol and | Statewide† | 8.6% | 16.6% 🔺 | 18.8% ▲ Above 75th |
| Other Drug Abuse Dependencies (FUA), 7-Day Follow-Up, Total | AMG | 12.2% | 15.5% | 18.1% Above 75th |
| | CCW | 15.4% | 12.1% | 13.4% Above 75th |
| | CHPW | 18.7% | 20.8% | 21.8% Above 75th |
| | MHW | 7.9% | 17.7% | 20.4% ▲ Above 75th |
| | UHC | NR | 12.2% | 16.4% ▲ Above 75th |
| ollow-Up After High Intensity Care for Substance Use Disorder | Statewide† | NR | 48.4% | 44.2% At 50th |
| FUI): 30-Day Follow-Up, 13-17 Years | AMG | NR | *** | *** At 50th |
| | CCW | NR | *** | *** At 50th |
| | CHPW | NR | *** | *** At 50th |
| | MHW | NR | *** | *** At 50th |
| | UHC | NR | *** | *** At 50th |
| ollow-Up After High Intensity Care for Substance Use Disorder | Statewide† | NR | 56.5% | 58.0% Above 50th, Below 75th |
| FUI): 30-Day Follow-Up, 18-64 Years | AMG | NR | 54.3% | 56.4% At 50th |
| • | CCW | NR | 51.2% | 55.8% At 50th |

| | CHPW | NR | 57.9% | 59.9% | At 75th |
|--|------------|----|-------|-------|------------------------|
| | MHW | NR | 58.8% | 59.0% | Above 50th, Below 75th |
| | UHC | NR | 51.8% | 56.7% | At 50th |
| Follow-Up After High Intensity Care for Substance Use Disorder | Statewide† | NR | *** | *** | At 50th |
| (FUI): 30-Day Follow-Up, 65+ Years | AMG | NR | *** | *** | At 50th |
| | CCW | NR | NR | *** | At 50th |
| | CHPW | NR | *** | NR | Below 50th |
| | MHW | NR | NR | *** | At 50th |
| | UHC | NR | NR | *** | At 50th |
| Follow-Up After High Intensity Care for Substance Use Disorder | Statewide† | NR | 56.5% | 57.9% | Above 50th, Below 75th |
| (FUI): 30-Day Follow-Up, Total | AMG | NR | 54.3% | 56.5% | Above 50th, Below 75th |
| | CCW | NR | 50.9% | 55.7% | At 50th |
| | CHPW | NR | 58.1% | 59.8% | At 75th |
| | MHW | NR | 58.7% | 58.9% | Above 50th, Below 75th |
| | UHC | NR | 51.9% | 56.4% | Above 50th, Below 75th |
| Follow-Up After High Intensity Care for Substance Use Disorder | Statewide† | NR | 35.5% | 28.8% | At 50th |
| (FUI): 7-Day Follow-Up, 13-17 Years | AMG | NR | *** | *** | Above 75th |
| | CCW | NR | *** | *** | At 75th |
| | CHPW | NR | *** | *** | At 50th |
| | MHW | NR | *** | *** | At 50th |
| | UHC | NR | *** | *** | At 50th |
| Follow-Up After High Intensity Care for Substance Use Disorder | Statewide† | NR | 39.0% | 38.3% | Above 50th, Below 75th |
| (FUI): 7-Day Follow-Up, 18-64 Years | AMG | NR | 40.5% | 37.9% | Above 50th, Below 75th |
| | CCW | NR | 37.4% | 35.5% | At 50th |
| | CHPW | NR | 36.0% | 41.0% | At 75th |
| | MHW | NR | 40.8% | 38.9% | Above 50th, Below 75th |
| | UHC | NR | 33.5% | 36.2% | Above 50th, Below 75th |
| Follow-Up After High Intensity Care for Substance Use Disorder | Statewide† | NR | *** | *** | At 50th |
| (FUI): 7-Day Follow-Up, 65+ Years | AMG | NR | *** | *** | At 50th |
| | CCW | NR | NR | *** | At 50th |
| | CHPW | NR | *** | NR | Below 50th |
| | MHW | NR | NR | *** | At 50th |
| | UHC | NR | NR | *** | At 50th |
| Follow-Up After High Intensity Care for Substance Use Disorder | Statewide† | NR | 39.0% | 38.2% | Above 50th, Below 75th |

| -UI): 7-Day Follow-Up, Total | AMG | NR | 40.5% | 38.1% | Above 50th, Below 75th |
|---|-------------|----------------|----------------|----------------|------------------------|
| (101). 7 Day 10110W Op, 10tul | CCW | NR | 37.3% | 35.6% | At 50th |
| | CHPW | NR | 36.3% | 40.9% | At 75th |
| | | | | | |
| | MHW | NR | 40.8% | 38.9% | Above 50th, Below 75th |
| harmanatharran fan Oriaid Haa Bisandar (BOD): 46 CA Varra | UHC | NR | 33.2% | 35.9% | Above 50th, Below 75th |
| harmacotherapy for Opioid Use Disorder (POD): 16-64 Years | Statewide† | NR | 18.9% | 19.4% | Below 50th |
| | AMG | NR | 18.4% | 21.0% | Below 50th |
| | CCW | NR | 19.3% | 17.6% | Below 50th |
| | CHPW | NR | 18.5% | 16.0% | Below 50th |
| | MHW | NR | 17.2% | 19.6% 🔺 | Below 50th |
| | UHC | NR | 24.3% | 21.5% | Below 50th |
| harmacotherapy for Opioid Use Disorder (POD): 65+ Years | Statewide† | NR | *** | 36.1% | At 50th |
| | AMG | NR | *** | *** | At 50th |
| | CCW | NR | *** | *** | At 50th |
| | CHPW | NR | *** | *** | At 50th |
| | MHW | NR | *** | *** | At 50th |
| | UHC | NR | *** | *** | At 50th |
| harmacotherapy for Opioid Use Disorder (POD): Total | Statewide† | NR | 18.9% | 19.5% | Below 50th |
| | AMG | NR | 18.4% | 21.0% | Below 50th |
| | CCW | NR | 19.4% | 17.7% | Below 50th |
| | CHPW | NR | 18.5% | 16.0% | Below 50th |
| | MHW | NR | 17.2% | 19.6% 🔺 | Below 50th |
| | UHC | NR | 24.5% | 21.6% | Below 50th |
| iabetes Screening for People With Schizophrenia or Bipolar | Statewide† | 84.6% | 82.0% 🔻 | 77.0% 🔻 | At 50th |
| isorder Who Are Using Antipsychotic Medication (SSD) | AMG | 86.8% | 83.6% | 76.8% 🔻 | At 50th |
| | CCW | 84.2% | 83.0% | 79.3% | At 75th |
| | CHPW | 85.6% | 79.9% ▼ | 73.9% 🔻 | Below 50th |
| | MHW | 83.4% | 82.4% | 78.2% ▼ | Above 50th, Below 75th |
| | UHC | 85.0% | 81.4% | 75.4% 🔻 | |
| iabetes Monitoring for People With Diabetes and Schizophrenia | | 65.4% | 67.5% | 60.9% ▼ | Below 50th |
| SMD) | AMG | 67.6% | 69.2% | 63.1% | At 50th |
| (SIVID) | AIVIU | | | | |
| | | | 63.6% | 59.2% | |
| | CCW CHPW | 65.3% 63.8% | 63.6% 60.3% | 59.2% 57.5% | At 50th Below 50th |

| | UHC | 66.9% | 72.1% | 61.4% | At 50th |
|---|--------------|-------|-------|-------|------------------------|
| Cardiovascular Monitoring for People with Cardiovascular Diseas | e Statewide† | 59.5% | 54.2% | 52.3% | Below 50th |
| and Schizophrenia (SMC) | AMG | *** | *** | *** | Below 50th |
| | CCW | *** | *** | *** | At 50th |
| | CHPW | *** | *** | *** | Below 50th |
| | MHW | 68.4% | 57.6% | 62.2% | At 50th |
| | UHC | *** | *** | *** | Below 50th |
| Adherence to Antipsychotic Medications for Individuals With | Statewide† | 63.8% | 65.5% | 67.1% | Above 50th, Below 75th |
| Schizophrenia (SAA) | AMG | 60.7% | 64.9% | 60.2% | Below 50th |
| | CCW | 60.8% | 61.6% | 64.1% | At 50th |
| | CHPW | 64.1% | 67.4% | 68.4% | At 75th |
| | MHW | 64.4% | 65.7% | 69.6% | At 75th |
| | UHC | 67.1% | 66.2% | 67.4% | At 75th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | NR | 42.6% | 36.4% | At 50th |
| Antipsychotics (APM), Blood Glucose Testing, 1-11 Years | AMG | NR | 32.9% | 32.9% | At 50th |
| | CCW | NR | 44.0% | 38.9% | At 50th |
| | CHPW | NR | 32.9% | 34.0% | At 50th |
| | MHW | NR | 44.9% | 37.1% | At 50th |
| | UHC | NR | 45.8% | 28.9% | At 50th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | NR | 59.0% | 52.8% | At 50th |
| Antipsychotics (APM), Blood Glucose Testing, 12-17 Years | AMG | NR | 65.7% | 54.7% | At 50th |
| | CCW | NR | 58.6% | 54.2% | At 50th |
| | CHPW | NR | 58.7% | 51.1% | At 50th |
| | MHW | NR | 59.8% | 52.2% | At 50th |
| | UHC | NR | 48.7% | 52.6% | At 50th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | NR | 53.3% | 47.5% | At 50th |
| Antipsychotics (APM), Blood Glucose Testing, Total | AMG | NR | 55.6% | 48.5% | At 50th |
| | CCW | NR | 53.5% | 49.1% | At 50th |
| | CHPW | NR | 50.0% | 45.8% | At 50th |
| | MHW | NR | 54.3% | 47.2% | At 50th |
| | UHC | NR | 47.8% | 45.5% | At 50th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | NR | 31.3% | 27.7% | At 50th |
| Antipsychotics (APM), Cholesterol Testing, 1-11 Years | AMG | NR | 20.3% | 28.2% | At 50th |
| | CCW | NR | 35.3% | 29.3% | At 50th |

| | CHPW | NR | 15.8% | 23.6% At 50th |
|--|------------|-------|---------|--------------------|
| | MHW | NR | 34.0% | 28.2% At 50th |
| | UHC | NR | 33.3% | 24.1% At 50th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | NR | 35.4% | 28.2% ▼ Below 50th |
| Antipsychotics (APM), Cholesterol Testing, 12-17 Years | AMG | NR | 39.3% | 32.7% At 50th |
| | CCW | NR | 37.2% | 30.2% At 50th |
| | CHPW | NR | 28.0% | 27.2% Below 50th |
| | MHW | NR | 37.0% | 26.9% ▼ Below 50th |
| | UHC | NR | 26.6% | 27.6% At 50th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | NR | 34.0% | 28.1% ▼ Below 50th |
| Antipsychotics (APM), Cholesterol Testing, Total | AMG | NR | 33.5% | 31.4% At 50th |
| | CCW | NR | 36.6% | 29.9% ▼ At 50th |
| | CHPW | NR | 23.8% | 26.1% Below 50th |
| | MHW | NR | 35.9% | 27.3% ▼ Below 50th |
| | UHC | NR | 28.7% | 26.5% At 50th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | 26.8% | 28.8% | 25.7% At 50th |
| Antipsychotics (APM), Blood Glucose and Cholesterol Testing, 1 | - AMG | 15.5% | 19.0% | 27.1% At 50th |
| 11 Years | CCW | 34.4% | 33.3% | 27.9% At 50th |
| | CHPW | 22.8% | 15.8% | 23.6% At 50th |
| | MHW | 25.7% | 30.4% | 25.8% At 50th |
| | UHC | 22.5% | 33.3% | 19.3% At 50th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | 30.2% | 33.9% | 27.5% ▼ Below 50th |
| Antipsychotics (APM), Blood Glucose and Cholesterol Testing, 1 | 2- AMG | 29.0% | 38.8% | 32.7% At 50th |
| 17 Years | CCW | 35.2% | 35.6% | 29.5% At 50th |
| | CHPW | 27.3% | 26.6% | 26.8% At 50th |
| | MHW | 29.3% | 35.3% 🔺 | 26.2% ▼ Below 50th |
| | UHC | 27.2% | 25.3% | 25.5% Below 50th |
| Metabolic Monitoring for Children and Adolescents on | Statewide† | 29.0% | 32.1% 🔺 | 26.9% ▼ Below 50th |
| Antipsychotics (APM), Blood Glucose and Cholesterol Testing, | AMG | 24.7% | 32.7% | 31.1% At 50th |
| Total | CCW | 34.9% | 34.8% | 29.0% At 50th |
| | CHPW | 25.9% | 22.9% | 25.8% At 50th |
| | MHW | 28.0% | 33.5% 🔺 | 26.1% ▼ Below 50th |
| | UHC | 25.8% | 27.8% | 23.7% Below 50th |
| Non-Recommended Cervical Cancer Screening in Adolescent | Statewide† | 0.5% | 0.3% 🔺 | 0.3% Above 75th |

| | AMG | 0.1% | 0.2% | 0.1% | Above 75th |
|--|------------|------|--------|----------|------------------------|
| | CCW | 0.4% | 0.4% | 0.4% | Above 50th, Below 75th |
| | CHPW | 0.4% | 0.3% | 0.2% | Above 75th |
| | MHW | 0.6% | 0.4% 🔺 | 0.3% | Above 50th, Below 75th |
| | UHC | 0.4% | 0.3% | 0.2% | Above 75th |
| Appropriate Treatment for Upper Respiratory Infection (URI), 3 | Statewide† | NR | 95.4% | 95.7% 🔺 | Above 75th |
| Months-17 Years | AMG | NR | 95.1% | 95.6% | Above 75th |
| | CCW | NR | 95.1% | 96.4% 🔺 | Above 75th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 95.6% | 96.0% | Above 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 95.4% | 95.3% | Above 75th |
| | UHC | NR | 95.4% | 96.0% | Above 75th |
| Appropriate Treatment for Upper Respiratory Infection (URI), 18- | Statewide† | NR | 89.0% | 90.8% | Above 75th |
| 64 Years | AMG | NR | 89.7% | 90.7% | Above 75th |
| | CCW | NR | 89.9% | 93.3% 🔺 | Above 75th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 89.2% | 91.7% 🔺 | Above 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 89.0% | 90.1% | Above 75th |
| | UHC | NR | 87.2% | 90.0% 🔺 | Above 75th |
| Appropriate Treatment for Upper Respiratory Infection (URI), 65+ | Statewide† | NR | 83.9% | 91.3% | Above 75th |
| Years | AMG | NR | *** | *** | At 50th |
| | CCW | NR | *** | *** | At 50th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | *** | *** | At 50th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | *** | *** | At 75th |
| | UHC | NR | *** | 90.3% 🔺 | At 75th |
| Appropriate Treatment for Upper Respiratory Infection (URI), | Statewide† | NR | 93.8% | 94.4% 🔺 | Above 75th |
| | AMG | NR | 93.4% | 94.1% | Above 75th |
| | CCW | NR | 94.1% | 95.7% 🔺 | Above 75th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 94.0% | 94.8% 🔺 | Above 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 94.0% | 94.0% | Above 75th |
| | UHC | NR | 93.0% | 94.0% 🔺 | Above 75th |
| Avoidance of Antibiotic Treatment for Acute | Statewide† | NR | 73.1% | 75.2% 🔺 | Above 75th |
| Bronchitis/Bronchiolitis (AAB), 3 Months-17 Years | AMG | NR | 79.8% | 79.0% | Above 75th |
| | CCW | NR | 80.7% | 82.1% | Above 75th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 72.5% | 73.2% | At 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 69.3% | 72.20/ 🛕 | At 75th |

| | UHC | NR | 76.2% | 79.0% Above 75th |
|---|------------|-------|---------|-------------------------------|
| Avoidance of Antibiotic Treatment for Acute | Statewide† | NR | 47.3% | 50.9% ▲ Above 75th |
| Bronchitis/Bronchiolitis (AAB), 18-64 Years | AMG | NR | 50.1% | 53.3% Above 75th |
| | CCW | NR | 50.9% | 54.5% Above 75th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 47.9% | 51.3% Above 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 45.7% | 48.9% ▲ Above 75th |
| | UHC | NR | 46.8% | 52.8% 🛕 Above 75th |
| Avoidance of Antibiotic Treatment for Acute | Statewide† | NR | *** | 26.2% Below 50th |
| Bronchitis/Bronchiolitis (AAB), 65+ Years | AMG | NR | *** | *** At 50th |
| | CCW | NR | NR | *** At 50th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | *** | *** At 50th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | *** | *** At 50th |
| | UHC | NR | *** | 22.9% Below 50th |
| Avoidance of Antibiotic Treatment for Acute | Statewide† | NR | 61.9% | 64.5% ▲ Above 75th |
| Bronchitis/Bronchiolitis (AAB), Total | AMG | NR | 64.1% | 66.3% Above 75th |
| | CCW | NR | 70.3% | 72.4% Above 75th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 61.7% | 63.6% At 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 59.5% | 62.4% 🔺 At 75th |
| | UHC | NR | 60.5% | 64.4% 🔺 Above 75th |
| Use of Imaging Studies for Low Back Pain (LBP) | Statewide† | 74.8% | 76.6% 🔺 | 76.9% Above 50th, Below 75th |
| | AMG | 74.1% | 77.6% | 77.8% At 75th |
| | CCW | 75.1% | 79.9% 🔺 | 76.6% At 50th |
| | CHPW | 76.1% | 76.5% | 80.6% 🔺 At 75th |
| | MHW | 74.3% | 75.3% | 75.6% At 50th |
| | UHC | 75.3% | 78.2% | 77.7% At 75th |
| Use of Opioids at High Dosage (HDO) (Note that a lower score is | Statewide† | NR | 7.1% | 6.3% ▲ Below 50th |
| better for this measure) | AMG | NR | 6.2% | 5.8% Above 50th, Below 75th |
| | CCW | NR | 7.1% | 6.5% Below 50th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 7.8% | 5.8% 🔺 Above 50th, Below 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 6.4% | 5.7% Below 50th |
| | UHC | NR | 9.4% | 8.3% Below 50th |
| Use of Opioids from Multiple Prescribers (UOP) (Note that a | Statewide† | 24.2% | 23.9% | 22.2% ▲ Below 50th |
| lower score is better for this measure) | AMG | 23.0% | 22.7% | 21.2% Below 50th |
| | CCW | 22.1% | 21.1% | 19.7% Above 50th, Below 75th |

| | CHPW | 25.5% | 23.7% | 21.4% Below 50th |
|---|--------------|-------|---------------|-------------------------------|
| | MHW | 25.0% | 25.1% | 23.3% A Below 50th |
| | UHC | 22.8% | 22.8% | 21.7% Below 50th |
| Use of Opioids from Multiple Pharmacies (UOP) (Note that a | Statewide† | 6.5% | 5.2% 🔺 | 3.7% ▲ Below 50th |
| ower score is better for this measure) | AMG | 4.8% | 3.4% | 2.8% Above 50th, Below 75th |
| | CCW | 6.6% | 5.2% | 3.4% 🔺 Above 50th, Below 75tl |
| | CHPW | 5.9% | 4.7% | 3.2% 🔺 Above 50th, Below 75tl |
| | MHW | 5.9% | 5.3% | 3.9% ▲ Below 50th |
| | UHC | 9.3% | 6.7% | 4.5% ▲ Below 50th |
| Jse of Opioids from Multiple Prescribers and Multiple Pharmacie | s Statewide† | 3.9% | 3.3% 🔺 | 2.5% ▲ Below 50th |
| UOP) (Note that a lower score is better for this measure) | AMG | 3.1% | 2.1% | 1.9% Above 50th, Below 75th |
| | CCW | 3.6% | 3.1% | 2.1% Above 50th, Below 75th |
| | CHPW | 3.8% | 3.1% | 2.0% 🔺 Above 50th, Below 75tl |
| | MHW | 3.8% | 3.5% | 2.7% ▲ Below 50th |
| | UHC | 5.0% | 3.9% 🔺 | 2.9% ▲ Below 50th |
| Risk of Continued Opioid Use (COU), At least 15 days, 18-64 Years | Statewide† | 7.3% | 6.7% ▼ | 6.3% ▼ At 50th |
| | AMG | 8.2% | 7.6% | 6.6% At 50th |
| | CCW | 6.1% | 5.7% | 5.8% At 50th |
| | CHPW | 7.3% | 7.1% | 6.4% At 50th |
| | MHW | 7.6% | 6.8% ▼ | 6.1% ▼ At 50th |
| | UHC | 7.0% | 6.3% | 7.0% Above 75th |
| Risk of Continued Opioid Use (COU), At least 15 days, 65+ Years | Statewide† | 13.1% | 5.4% | 19.7% ▲ Above 75th |
| | AMG | *** | *** | *** At 50th |
| | CCW | *** | *** | *** At 50th |
| | CHPW | *** | *** | *** Below 50th |
| | MHW | 14.6% | 5.0% | *** At 50th |
| | UHC | *** | *** | 21.3% Above 75th |
| Risk of Continued Opioid Use (COU), At least 15 days, Total | Statewide† | 7.3% | 6.7% ▼ | 6.4% At 50th |
| | AMG | 8.2% | 7.6% | 6.6% ▼ At 50th |
| | CCW | 6.1% | 5.7% | 5.8% Below 50th |
| | CHPW | 7.3% | 7.1% | 6.4% At 50th |
| | MHW | 7.6% | 6.8% ▼ | 6.1% ▼ At 50th |
| | UHC | 7.0% | 6.3% | 7.6% 🔺 Above 75th |
| Risk of Continued Opioid Use (COU), At least 30 days, 18-64 Years | Statewide† | 2.7% | 2.8% | 2.7% Below 50th |

| | AMG | 3.3% | 3.5% | 2.9% Be | low 50th |
|---|------------|-------|---------|------------|----------|
| | CCW | 2.1% | 2.3% | | low 50th |
| | CHPW | 2.6% | 3.1% | | low 50th |
| | MHW | 2.9% | 2.8% | | low 50th |
| | UHC | 2.6% | 2.4% | 3.2% ▲ At | 50th |
| Risk of Continued Opioid Use (COU), At least 30 days, 65+ Years | Statewide† | 2.5% | 3.2% | 11.2% At | 50th |
| | AMG | *** | *** | *** At | 50th |
| | CCW | *** | *** | *** At | 50th |
| | CHPW | *** | *** | *** At | 50th |
| | MHW | 2.4% | 2.5% | *** At | 50th |
| | UHC | *** | *** | 12.8% Ab | ove 75th |
| Risk of Continued Opioid Use (COU), At least 30 days, Total | Statewide† | 2.7% | 2.8% | 2.7% Be | low 50th |
| | AMG | 3.3% | 3.5% | 2.9% Be | low 50th |
| | CCW | 2.1% | 2.3% | 2.1% Be | low 50th |
| | CHPW | 2.6% | 3.1% | 2.5% Be | low 50th |
| | MHW | 2.9% | 2.8% | 2.6% Be | low 50th |
| | UHC | 2.6% | 2.4% | 3.6% ▲ At | 50th |
| Adults' Access to Preventive/Ambulatory Health Services (AAP), | Statewide† | 73.1% | 74.1% 🔺 | 70.9% ▼ Be | low 50th |
| 20-44 Years | AMG | 66.8% | 69.8% 🔺 | 65.7% ▼ Be | low 50th |
| | CCW | 69.0% | 70.4% 🔺 | 68.7% ▼ Be | low 50th |
| | CHPW | 72.0% | 73.1% 🔺 | 70.5% ▼ Be | low 50th |
| | MHW | 77.5% | 77.5% | 73.8% ▼ Be | low 50th |
| | UHC | 70.1% | 71.2% | 68.2% ▼ Be | low 50th |
| Adults' Access to Preventive/Ambulatory Health Services (AAP), | Statewide† | 80.2% | 80.5% | 77.2% ▼ Be | low 50th |
| 45-64 Years | AMG | 75.1% | 76.8% 🔺 | 72.7% ▼ Be | low 50th |
| | CCW | 77.0% | 77.8% | 75.4% ▼ Be | low 50th |
| | CHPW | 80.5% | 80.4% | 77.7% 🔻 Be | low 50th |
| | MHW | 83.3% | 83.0% | 79.4% ▼ Be | low 50th |
| | UHC | 79.4% | 79.7% | 76.9% ▼ Be | low 50th |
| Adults' Access to Preventive/Ambulatory Health Services (AAP), | Statewide† | 65.0% | 94.1% 🔺 | 88.6% ▼ At | 75th |
| 55+ Years | AMG | 27.3% | 84.2% | 68.4% ▼ Be | low 50th |
| | CCW | 86.1% | 86.2% | 77.8% Be | low 50th |
| | CHPW | 88.3% | 89.4% | 79.8% ▼ At | |
| | CHPVV | 00.5% | 03.470 | 73.0% ▼ At | 50(1) |

| | UHC | 88.5% | 96.3% 🔺 | 94.0% | Above 75th |
|--|------------|-------|---------|-------|------------|
| Adults' Access to Preventive/Ambulatory Health Services (AAP), | Statewide† | 75.5% | 76.5% 🔺 | 73.2% | Below 50th |
| Total | AMG | 69.5% | 72.5% 🔺 | 68.2% | Below 50th |
| | CCW | 71.9% | 73.1% 🔺 | 71.0% | Below 50th |
| | CHPW | 75.2% | 75.9% 🔺 | 73.1% | Below 50th |
| | MHW | 79.3% | 79.3% | 75.5% | Below 50th |
| | UHC | 73.6% | 75.3% 🔺 | 72.4% | Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 29.7% | 25.8% | 32.6% | Below 50th |
| Dependence Treatment (IET), Alcohol Abuse or Dependence | AMG | *** | 23.5% | 32.6% | At 50th |
| Initiation of AOD Treatment: 13-17 Years | CCW | *** | 32.1% | 35.4% | At 50th |
| | CHPW | *** | 22.8% | 21.2% | Below 50th |
| | MHW | 30.5% | 23.4% | 34.4% | Below 50th |
| | UHC | NR | *** | 31.7% | At 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 11.9% | 6.9% | 8.8% | At 50th |
| Dependence Treatment (IET), Alcohol Abuse or Dependence | AMG | *** | 5.9% | 4.7% | At 50th |
| Engagement of AOD Treatment: 13-17 Years | CCW | *** | 11.0% | 10.6% | At 50th |
| | CHPW | *** | 3.5% | 6.1% | At 50th |
| | MHW | 14.7% | 5.9% | 9.5% | At 50th |
| | UHC | NR | *** | 7.3% | At 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | *** | 57.6% | 38.9% | At 50th |
| Dependence Treatment (IET), Opioid Abuse or Dependence: | AMG | NR | *** | *** | At 50th |
| Initiation of AOD Treatment: 13-17 Years | CCW | NR | *** | *** | At 50th |
| | CHPW | NR | *** | *** | At 50th |
| | MHW | *** | *** | *** | At 50th |
| | UHC | NR | *** | *** | At 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | *** | 15.2% | 14.8% | At 50th |
| Dependence Treatment (IET), Opioid Abuse or Dependence: | AMG | NR | *** | *** | At 50th |
| Engagement of AOD Treatment: 13-17 Years | CCW | NR | *** | *** | At 50th |
| | CHPW | NR | *** | *** | At 50th |
| | MHW | *** | *** | *** | At 50th |
| | UHC | NR | *** | *** | At 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 31.3% | 32.9% | 37.9% | Below 50th |
| Dependence Treatment (IET), Other Drug Abuse or Dependence: | AMG | *** | 41.2% | 43.2% | At 50th |
| Initiation of AOD Treatment: 13-17 Years | CCW | *** | 33.5% 🔺 | 38.9% | Below 50th |

| | CHPW | *** | 30.1% | 38.5% Below 50th |
|---|------------|-------|---------|--------------------------------|
| | MHW | 34.8% | 33.1% | 35.9% Below 50th |
| | UHC | NR | 20.8% | 40.0% At 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 17.2% | 10.8% ▼ | 11.7% Below 50th |
| Dependence Treatment (IET), Other Drug Abuse or Dependence: | AMG | *** | 14.7% | 12.9% At 50th |
| Engagement of AOD Treatment: 13-17 Years | CCW | *** | 10.9% | 15.4% At 50th |
| | CHPW | *** | 6.4% | 13.0% At 50th |
| | MHW | 19.9% | 11.6% 🔻 | 10.6% Below 50th |
| | UHC | NR | 5.7% | 6.7% Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 28.0% | 30.9% | 35.6% ▲ Below 50th |
| Dependence Treatment (IET), Total: Initiation of AOD Treatment: | AMG | *** | 36.4% | 41.9% At 50th |
| 13-17 Years | CCW | 12.5% | 32.2% | 36.5% Below 50th |
| | CHPW | *** | 28.4% | 34.0% Below 50th |
| | MHW | 30.0% | 30.6% | 34.4% Below 50th |
| | UHC | NR | 23.8% | 35.8% Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 13.8% | 9.7% | 10.7% Below 50th |
| Dependence Treatment (IET), Total: Engagement of AOD | AMG | *** | 12.4% | 11.3% At 50th |
| Treatment: 13-17 Years | CCW | 6.3% | 10.3% | 13.7% At 50th |
| | CHPW | *** | 5.7% | 10.5% At 50th |
| | MHW | 15.9% | 10.3% | 10.0% Below 50th |
| | UHC | NR | 6.3% | 7.3% At 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 31.9% | 35.4% 🔺 | 38.4% ▲ Below 50th |
| Dependence Treatment (IET), Alcohol Abuse or Dependence | AMG | 48.0% | 37.8% | 41.2% At 50th |
| Initiation of AOD Treatment: 18+ Years | CCW | 36.3% | 34.0% | 34.6% Below 50th |
| | CHPW | 37.5% | 29.6% 🔻 | 33.7% Below 50th |
| | MHW | 31.1% | 36.1% 🔺 | 39.7% A Below 50th |
| | UHC | NR | 37.9% | 38.4% Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 7.0% | 9.1% 🔺 | 11.4% 🔺 At 50th |
| Dependence Treatment (IET), Alcohol Abuse or Dependence | AMG | 22.0% | 9.2% | 12.0% 🔺 At 50th |
| Engagement of AOD Treatment: 18+ Years | CCW | 12.3% | 7.2% | 9.4% Below 50th |
| | CHPW | 11.0% | 7.5% | 11.5% 🔺 At 50th |
| | MHW | 6.4% | 10.0% 🔺 | 12.1% 🔺 At 50th |
| | UHC | NR | 8.5% | 9.7% Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 60.0% | 62.5% | 64.7% 🔺 Above 50th, Below 75th |

| Dependence Treatment (IET), Opioid Abuse or Dependence: | AMG | *** | 63.1% | 64.5% At 75th |
|---|------------|-------------|----------------|--------------------------------|
| Initiation of AOD Treatment: 18+ Years | CCW | 51.2% | 62.4% | 60.0% At 50th |
| middlen of Nob Treatment. 10. Tears | CHPW | 70.2% | 60.7% | 66.5% ▲ At 75th |
| | MHW | 59.8% | 63.2% | 65.7% At 75th |
| | UHC | NR | 59.2% | 63.4% Above 50th, Below 75th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 34.7% | 34.4% | 34.2% Above 50th, Below 75th |
| Dependence Treatment (IET), Opioid Abuse or Dependence: | AMG | *** | 33.5% | 32.3% Above 50th, Below 75th |
| Engagement of AOD Treatment: 18+ Years | CCW | 24.4% | 31.9% | 31.0% At 50th |
| Engagement of Nob Treatment. 10. Tears | CHPW | 37.6% | 33.2% | 35.0% Above 50th, Below 75th |
| | MHW | 34.8% | 36.0% | 36.0% Above 50th, Below 75th |
| | UHC | 94.8% NR | 29.8% | 32.5% Above 50th, Below 75th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 32.5% | 36.4% 🔺 | 38.3% ▲ Below 50th |
| Dependence Treatment (IET), Other Drug Abuse or Dependence: | AMG | 31.8% | 40.8% | 43.0% At 50th |
| Initiation of AOD Treatment: 18+ Years | CCW | 31.4% | 34.8% | 35.4% Below 50th |
| initiation of AOD Treatment. 101 Tears | CHPW | 41.1% | 29.2% ▼ | 31.2% Below 50th |
| | MHW | 32.1% | 37.1% ▲ | 39.2% ▲ Below 50th |
| | UHC | 32.1% NR | 37.1% | 38.8% Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 8.2% | 9.1% | 9.5% Below 50th |
| Dependence Treatment (IET), Other Drug Abuse or Dependence: | AMG | 9.1% | 9.1% | 9.4% Below 50th |
| Engagement of AOD Treatment: 18+ Years | CCW | 8.7% | 8.0% | 7.4% Below 50th |
| Engagement of AOD Treatment. 101 Tears | CHPW | 12.6% | 10.0% | 10.1% At 50th |
| | MHW | 8.0% | 9.5% | 10.0% Below 50th |
| | UHC | NR | 6.3% | 9.3% A Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 38.8% | 42.5% | 45.4% Above 50th, Below 75th |
| Dependence Treatment (IET), Total: Initiation of AOD Treatment: | | 40.5% | 45.4% | 47.8% At 75th |
| 18+ Years | CCW | 36.8% | 40.2% | 40.9% Below 50th |
| 10+ 16013 | CHPW | 46.5% | 38.6% ▼ | 42.9% ▲ Below 50th |
| | MHW | 38.5% | 43.1% ▲ | 46.5% ▲ Above 50th, Below 75th |
| | UHC | 36.3% NR | 43.3% | 45.2% At 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 14.6% | 15.1% | 16.6% A Above 50th, Below 75th |
| Dependence Treatment (IET), Total: Engagement of AOD | AMG | 14.6% | 15.1% 15.2% | 16.1% Above 50th, Below 75th |
| Treatment: 18+ Years | CCW | 17.6% | | 13.9% At 50th |
| Treatment: 18+ feats | | | 13.1% | |
| | CHPW | 16.6% | 13.3% | 16.4% A Above 50th, Below 75th |
| | MHW | 14.5% | 16.2% | 17.6% 🔺 Above 50th, Below 75th |

| | UHC | NR | 13.3% | 15.9% 🛕 Above 50th, Below 75th |
|---|------------|-------|---------|--------------------------------|
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 31.8% | 35.1% 🔺 | 38.2% 🛕 Below 50th |
| Dependence Treatment (IET), Alcohol Abuse or Dependence: | AMG | 47.1% | 37.5% | 41.0% Below 50th |
| Initiation of AOD Treatment: Total | CCW | 35.1% | 33.8% | 34.6% Below 50th |
| | CHPW | 37.4% | 29.4% 🔻 | 33.3% Below 50th |
| | MHW | 31.1% | 35.7% 🔺 | 39.5% 🛕 Below 50th |
| | UHC | NR | 37.9% | 38.3% Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 7.1% | 9.1% 🔺 | 11.3% 🔺 At 50th |
| Dependence Treatment (IET), Alcohol Abuse or Dependence: | AMG | 21.6% | 9.2% 🔻 | 11.9% 🔺 At 50th |
| Engagement of AOD Treatment: Total | CCW | 11.7% | 7.5% | 9.4% Below 50th |
| | CHPW | 10.5% | 7.3% | 11.4% 🔺 At 50th |
| | MHW | 6.6% | 9.9% 🔺 | 12.0% 🔺 Above 50th, Below 75th |
| | UHC | NR | 8.5% | 9.7% Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 60.0% | 62.4% | 64.6% 🔺 Above 50th, Below 75th |
| Dependence Treatment (IET), Opioid Abuse or Dependence: | AMG | *** | 63.0% | 64.5% At 75th |
| Initiation of AOD Treatment: Total | CCW | 51.2% | 62.4% | 59.7% At 50th |
| | CHPW | 70.2% | 60.5% | 66.4% 🔺 At 75th |
| | MHW | 59.8% | 63.3% 🔺 | 65.6% At 75th |
| | UHC | NR | 59.1% | 63.4% Above 50th, Below 75th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 34.6% | 34.3% | 34.1% Above 50th, Below 75th |
| Dependence Treatment (IET), Opioid Abuse or Dependence: | AMG | *** | 33.5% | 32.3% Above 50th, Below 75th |
| Engagement of AOD Treatment: Total | CCW | 24.4% | 31.7% | 30.8% At 50th |
| | CHPW | 37.6% | 33.1% | 34.8% Above 50th, Below 75th |
| | MHW | 34.7% | 36.0% | 35.9% Above 50th, Below 75th |
| | UHC | NR | 29.7% | 32.5% Above 50th, Below 75th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 32.5% | 36.1% 🔺 | 38.3% ▲ Below 50th |
| Dependence Treatment (IET), Other Drug Abuse or Dependence: | AMG | 30.4% | 40.9% | 43.0% At 50th |
| Initiation of AOD Treatment: Total | CCW | 28.5% | 34.6% | 35.7% Below 50th |
| | CHPW | 40.7% | 29.3% 🔻 | 31.6% Below 50th |
| | MHW | 32.2% | 36.8% 🔺 | 39.0% 🛕 Below 50th |
| | UHC | NR | 37.2% | 38.8% Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 8.5% | 9.2% | 9.6% Below 50th |
| Dependence Treatment (IET), Other Drug Abuse or Dependence: | AMG | 8.7% | 9.4% | 9.5% Below 50th |
| Engagement of AOD Treatment: Total | CCW | 8.5% | 8.4% | 8.1% Below 50th |

| | CHPW | 12.3% | 9.7% | 10.2% | At 50th |
|--|------------|-------|----------------|----------------|------------------------|
| | MHW | 8.3% | 9.7% 🔺 | 10.1% | Below 50th |
| | UHC | NR | 6.2% | 9.2% 🔺 | Below 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 38.6% | 42.0% 🔺 | 45.1% 🔺 | At 50th |
| Dependence Treatment (IET), Total: Initiation of AOD Treatment: | AMG | 39.6% | 45.2% | 47.6% | Above 50th, Below 75th |
| Total | CCW | 34.8% | 39.5% | 40.7% | Below 50th |
| | CHPW | 46.0% | 38.1% T | 42.5% | Below 50th |
| | MHW | 38.3% | 42.5% | 46.0% 🔺 | Above 50th, Below 75th |
| | UHC | NR | 42.9% | 45.0% | At 50th |
| Initiation and Engagement of Alcohol and Other Drug | Statewide† | 14.6% | 14.8% | 16.3% 🔺 | Above 50th, Below 75th |
| Dependence Treatment (IET), Total: Engagement of AOD | AMG | 17.2% | 15.1% | 16.0% | Above 50th, Below 75th |
| Treatment: Total | CCW | 13.1% | 12.8% | 13.9% | At 50th |
| | CHPW | 16.2% | 12.9% | 16.2% 🔺 | Above 50th, Below 75th |
| | MHW | 14.5% | 15.9% 🔺 | 17.3% 🔺 | At 75th |
| | UHC | NR | 13.1% | 15.7% 🔺 | Above 50th, Below 75th |
| Prenatal and Postpartum Care (PPC), Timeliness of Prenatal Care | Statewide† | NR | 87.2% | 82.7% | Below 50th |
| | AMG | NR | 76.3% | 78.6% | Below 50th |
| | CCW | NR | 77.4% | 78.1% | Below 50th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 76.4% | 88.3% 🔺 | At 50th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 94.6% | 82.5% ▼ | Below 50th |
| | UHC | NR | 88.3% | 86.1% | At 50th |
| Prenatal and Postpartum Care (PPC), Postpartum Care | Statewide† | NR | 73.6% | 76.7% 🔺 | At 50th |
| | AMG | NR | 60.1% | 71.8% 🔺 | Below 50th |
| | CCW | NR | 72.7% | 73.5% | At 50th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 77.4% | 82.5% | At 75th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 75.2% | 77.4% | At 50th |
| | UHC | NR | 74.7% | 75.2% | At 50th |
| Use of First-Line Psychosocial Care for Children and Adolescents | Statewide† | 12.7% | 55.8% 🔺 | 62.9% | At 50th |
| on Antipsychotics (APP), 1-11 Years | AMG | *** | 40.6% 🔺 | 50.0% | At 50th |
| | CCW | 9.0% | 52.9% 🔺 | 62.4% | At 50th |
| | CHPW | 6.6% | 45.9% 🔺 | *** | At 50th |
| | MHW | 57.9% | 67.6% | 66.5% | At 50th |
| | UHC | 2.0% | *** | *** | At 50th |
| Use of First-Line Psychosocial Care for Children and Adolescents | Statewide† | 18.3% | 55.7% 🔺 | 60.4% | At 50th |

| on Antipsychotics (APP), 12-17 Years | AMG | 9.1% | 34.5% 🔺 | 52.4% | Below 50th |
|--|------------|-------|---------|-------|------------|
| | CCW | 13.2% | 55.6% 🔺 | 66.0% | At 50th |
| | CHPW | 13.2% | 50.0% 🔺 | 54.8% | At 50th |
| | MHW | 61.7% | 69.0% | 63.1% | At 50th |
| | UHC | 6.3% | 43.1% 🔺 | 49.3% | Below 50th |
| Use of First-Line Psychosocial Care for Children and Adolescents | Statewide† | 16.6% | 55.7% 🔺 | 61.2% | At 50th |
| on Antipsychotics (APP), Total | AMG | 7.8% | 36.7% | 51.7% | Below 50th |
| | CCW | 11.7% | 54.7% | 64.7% | At 50th |
| | CHPW | 11.3% | 48.6% | 55.6% | At 50th |
| | MHW | 60.7% | 68.5% | 64.2% | At 50th |
| | UHC | 5.2% | 47.0% | 52.2% | Below 50th |
| Well-Child Visits in the First 30 Months of Life (W30), 0-15 | Statewide† | NR | NR | 54.0% | Below 50th |
| Months | AMG | NR | NR | 49.5% | Below 50th |
| | CCW | NR | NR | 55.7% | At 50th |
| | CHPW | NR | NR | 62.2% | At 75th |
| | MHW | NR | NR | 54.3% | At 50th |
| | UHC | NR | NR | 44.9% | Below 50th |
| Well-Child Visits in the First 30 Months of Life (W30), 16-30 | Statewide† | NR | NR | 68.4% | Below 50th |
| Months | AMG | NR | NR | 66.9% | Below 50th |
| | CCW | NR | NR | 71.8% | At 50th |
| | CHPW | NR | NR | 67.3% | Below 50th |
| | MHW | NR | NR | 68.3% | Below 50th |
| | UHC | NR | NR | 67.4% | Below 50th |
| Child and Adolescent Well-Care Visit (WCV), Age 3-11 | Statewide† | NR | NR | 46.9% | Below 50th |
| | AMG | NR | NR | 42.2% | Below 50th |
| | CCW | NR | NR | 49.3% | Below 50th |
| | CHPW | NR | NR | 44.9% | Below 50th |
| | MHW | NR | NR | 48.4% | Below 50th |
| | UHC | NR | NR | 43.3% | Below 50th |
| Child and Adolescent Well-Care Visit (WCV), Age 12-17 | Statewide† | NR | NR | 34.8% | Below 50th |
| | AMG | NR | NR | 29.3% | Below 50th |
| | CCW | NR | NR | 36.3% | Below 50th |
| | CHPW | NR | NR | 33.8% | Below 50th |
| | MHW | NR | NR | 36.2% | Below 50th |
| | | | | | |

| | UHC | NR | NR | 31.5% | Below 50th |
|--|------------|-------|--------|--------|------------------------|
| Child and Adolescent Well-Care Visit (WCV), Age 18-21 | Statewide† | NR | NR | 17.7% | Below 50th |
| | AMG | NR | NR | 14.6% | Below 50th |
| | CCW | NR | NR | 16.4% | Below 50th |
| | CHPW | NR | NR | 17.4% | Below 50th |
| | MHW | NR | NR | 19.0% | Below 50th |
| | UHC | NR | NR | 16.7% | Below 50th |
| Child and Adolescent Well-Care Visit (WCV), Total | Statewide† | NR | NR | 38.8% | Below 50th |
| | AMG | NR | NR | 34.0% | Below 50th |
| | CCW | NR | NR | 40.4% | Below 50th |
| | CHPW | NR | NR | 36.8% | Below 50th |
| | MHW | NR | NR | 40.3% | Below 50th |
| | UHC | NR | NR | 35.7% | Below 50th |
| Ambulatory Care (AMB), Outpatient | Statewide† | 301.1 | 304.1 | 251.5 | NR |
| | AMG | 252.6 | 260.0 | 212.6 | NR |
| | CCW | 304.1 | 307.2 | 257.8 | NR |
| | CHPW | 279.9 | 285.0 | 218.8 | NR |
| | MHW | 320.8 | 322.1 | 264.9 | NR |
| | UHC | 294.2 | 296.9 | 266.1 | NR |
| Ambulatory Care (AMB), Emergency Department | Statewide† | 48.1 | 48.3 | 35.8 | NR |
| | AMG | 50.9 | 50.6 | 38.9 | NR |
| | CCW | 53.0 | 53.9 | 37.4 | NR |
| | CHPW | 49.7 | 49.9 | 35.0 | NR |
| | MHW | 46.0 | 46.7 | 34.5 | NR |
| | UHC | 46.3 | 44.6 | 37.2 | NR |
| Identification of Alcohol and Other Drug Services (IAD), Total Any | Statewide† | 7.0% | 8.2% 🔺 | 8.1% | Above 50th, Below 75th |
| Services | AMG | 8.7% | 10.4% | 10.3% | Above 75th |
| | CCW | 6.4% | 7.6% 🔺 | 7.1% 🔻 | Above 50th, Below 75th |
| | CHPW | 11.0% | 8.3% | 7.8% | Above 50th, Below 75th |
| | MHW | 6.5% | 7.4% 🛕 | 7.6% 🔺 | Above 50th, Below 75th |
| | UHC | NR | 10.4% | 9.4% | At 75th |
| Identification of Alcohol and Other Drug Services (IAD), Total | Statewide† | 1.6% | 1.6% | 1.6% 🔺 | Above 50th, Below 75th |
| Inpatient | AMG | 1.2% | 2.1% 🔺 | 2.3% 🔺 | Above 75th |
| | CCW | 1.2% | 1.4% | 1.3% | Below 50th |

| | CHPW | 2.2% | 1.6% ▼ | 1.6% | Above 50th, Below 75th |
|--|------------|------|--------|-------|------------------------|
| | MHW | 1.6% | 1.4% ▼ | 1.5% | Below 50th |
| | UHC | NR | 1.9% | 1.8% | Above 50th, Below 75th |
| Identification of Alcohol and Other Drug Services (IAD), Total | Statewide† | 0.4% | 0.0% ▼ | 0.1% | Below 50th |
| Intensive Outpatient/Partial Hospitalization | AMG | 0.0% | 0.0% | 0.1% | Below 50th |
| | CCW | 0.0% | 0.0% | 0.0% | Below 50th |
| | CHPW | 0.0% | 0.0% | 0.1% | At 50th |
| | MHW | 0.5% | 0.0% | 0.1% | Above 50th, Below 75th |
| | UHC | NR | 0.1% | 0.1% | Below 50th |
| dentification of Alcohol and Other Drug Services (IAD), Total | Statewide† | 5.6% | 6.5% 🔺 | 6.2% | Above 50th, Below 75th |
| Outpatient/MAT | AMG | 6.8% | 7.9% 🛕 | 7.7% | Above 75th |
| | CCW | 5.3% | 5.8% | 5.2% | Above 50th, Below 75th |
| | CHPW | 8.8% | 6.6% ▼ | 6.0% | Above 50th, Below 75th |
| | MHW | 5.2% | 6.0% 🔺 | 5.9% | Above 50th, Below 75th |
| | UHC | NR | 8.4% | 7.3% | Above 75th |
| dentification of Alcohol and Other Drug Services (IAD), Total ED | Statewide† | 2.2% | 2.5% 🔺 | 2.3% | Above 50th, Below 75th |
| | AMG | 2.6% | 3.4% | 3.0% | Above 75th |
| | CCW | 1.8% | 2.5% | 2.3% | Above 50th, Below 75th |
| | CHPW | 2.9% | 2.5% 🔻 | 2.3% | Above 50th, Below 75th |
| | MHW | 2.1% | 2.1% | 2.1% | Above 50th, Below 75th |
| | UHC | NR | 2.8% | 2.5% | Above 50th, Below 75th |
| dentification of Alcohol and Other Drug Services (IAD), Total | Statewide† | 0.5% | 0.6% 🔺 | 2.6% | Above 50th, Below 75th |
| Telehealth | AMG | 0.1% | 0.8% | 2.9% | Above 50th, Below 75th |
| | CCW | 0.0% | 0.5% | 2.3% | Above 50th, Below 75th |
| | CHPW | 1.1% | 0.4% | 2.0% | Above 50th, Below 75th |
| | MHW | 0.4% | 0.6% | 2.6% | Above 50th, Below 75th |
| | UHC | NR | 0.6% | 2.8% | Above 50th, Below 75th |
| Mental Health Utilization (MPT), Total Outpatient | Statewide† | 6.5% | 11.2% | 10.6% | NR |
| | AMG | 4.6% | 7.6% | 7.3% | NR |
| | CCW | 6.6% | 12.0% | 10.5% | NR |
| | CHPW | 5.2% | 8.8% | 8.6% | NR |
| | MHW | 9.9% | 13.3% | 12.0% | NR |
| | UHC | 4.2% | 10.1% | 10.6% | NR |
| Mental Health Utilization (MPT), Total ED | Statewide† | 0.6% | 0.3% | 0.6% | NR |

| | AMG | 0.2% | 0.1% | 0.1% | NR |
|---|--------------|-------|---------|---------|-----------------|
| | CCW | 0.1% | 0.1% | 0.1% | NR |
| | CHPW | 0.1% | 0.1% | 0.1% | NR |
| | MHW | 1.8% | 0.5% | 1.2% | NR |
| | UHC | 0.2% | 0.1% | 0.1% | NR |
| Plan All-Cause Readmissions (PCR), Total: 18-44 Years | Statewide† | NR | 8.4% | 9.0% | No Benchmark |
| | AMG | NR | 8.6% | 9.1% | No Benchmark |
| | CCW | NR | 9.2% | 9.4% | No Benchmark |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 10.4% | 10.1% | No Benchmark |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 7.9% | 8.5% | No Benchmark |
| | UHC | NR | 7.5% | 9.9% | No Benchmark |
| Plan All-Cause Readmissions (PCR), Total: 45-54 Years | Statewide† | NR | 9.2% | 9.0% | No Benchmark |
| | AMG | NR | 9.2% | 9.6% | No Benchmark |
| | CCW | NR | 10.0% | 7.4% | No Benchmark |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 10.6% | 9.2% | No Benchmark |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 8.6% | 8.6% | No Benchmark |
| | UHC | NR | 8.9% | 10.5% | No Benchmark |
| Plan All-Cause Readmissions (PCR), Total: 55-64 Years | Statewide† | NR | 9.6% | 9.8% | No Benchmark |
| | AMG | NR | 8.1% | 9.9% | No Benchmark |
| | CCW | NR | 9.9% | 9.9% | No Benchmark |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 11.0% | 9.8% | No Benchmark |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 9.2% | 9.5% | No Benchmark |
| | UHC | NR | 10.5% | 10.6% | No Benchmark |
| Plan All-Cause Readmissions (PCR), Total | Statewide† | NR | 9.0% | 9.3% | Below 50th |
| | AMG | NR | 8.6% | 9.5% | Below 50th |
| | CCW | NR | 9.6% | 9.1% | Below 50th |
| Due to substantial changes in measure specifications between MY | CHPW | NR | 10.6% | 9.8% | Below 50th |
| 2018 and MY 2019, MY 2018 results are not displayed. | MHW | NR | 8.4% | 8.8% | Below 50th |
| | UHC | NR | 8.9% | 10.3% | Below 50th |
| Substance Use Disorder Treatment Penetration (SUD), 12-64 years | s Statewide† | 34.1% | 36.6% 🔺 | 38.4% 🔺 | At Benchmark |
| | AMG | 33.6% | 34.9% 🔺 | 38.9% 🔺 | At Benchmark |
| | CCW | 31.5% | 34.4% 🔺 | 35.2% | Below Benchmark |
| | CHPW | 35.4% | 38.2% 🔺 | 39.6% | Above Benchmark |
| | MHW | 34.4% | 37.8% | 38.5% | At Benchmark |

| | UHC | 34.5% | 35.5% | 39.0% ▲ Above Benchmark |
|---|-----------------|-------|----------------|-------------------------|
| Substance Use Disorder Treatment Penetration (SUD), 12-17 years | ears Statewide† | 33.0% | 28.2% 🔻 | 24.4% ▼ Below Benchmark |
| | AMG | 31.8% | 29.5% | 27.5% At Benchmark |
| | CCW | 36.5% | 27.9% 🔻 | 23.2% Below Benchmark |
| | CHPW | 35.2% | 27.4% 🔻 | 21.4% ▼ Below Benchmark |
| | MHW | 32.2% | 28.5% 🔻 | 25.1% ▼ Below Benchmark |
| | UHC | 26.4% | 26.6% | 23.6% At Benchmark |
| Substance Use Disorder Treatment Penetration (SUD), 12-26 | Statewide† | 29.6% | 29.3% | 27.7% ▼ Below Benchmark |
| years | AMG | 30.8% | 31.8% | 30.9% At Benchmark |
| | CCW | 29.5% | 28.6% | 25.8% Below Benchmark |
| | CHPW | 31.6% | 28.9% | 27.2% At Benchmark |
| | MHW | 29.0% | 29.3% | 27.3% ▼ Below Benchmark |
| | UHC | 28.9% | 28.6% | 29.0% At Benchmark |
| Substance Use Disorder Treatment Penetration (SUD), 18-64 ye | ears Statewide† | 34.2% | 37.6% 🔺 | 39.1% ▲ Above Benchmark |
| | AMG | 33.9% | 37.6% | 39.1% ▲ At Benchmark |
| | CCW | 31.2% | 35.0% 🔺 | 36.3% Below Benchmark |
| | CHPW | 35.4% | 38.4% 🔺 | 40.4% ▲ Above Benchmark |
| | MHW | 34.4% | 38.1% 🔺 | 39.1% ▲ Above Benchmark |
| | UHC | 34.9% | 37.0% 🔺 | 39.3% ▲ Above Benchmark |
| Mental Health Treatment Penetration (MH-B), 6-64 years | Statewide† | 57.3% | 57.4% | 53.9% ▼ Below Benchmark |
| | AMG | 63.7% | 62.6% ▼ | 51.5% ▼ Below Benchmark |
| | CCW | 54.6% | 54.3% | 55.4% ▲ Below Benchmark |
| | CHPW | 51.4% | 52.7% | 54.0% ▲ Below Benchmark |
| | MHW | 52.6% | 53.4% | 55.3% ▲ Below Benchmark |
| | UHC | 59.1% | 59.4% | 48.9% ▼ Below Benchmark |
| Mental Health Treatment Penetration (MH-B), 6-17 years | Statewide† | 67.9% | 66.4% ▼ | 64.0% ▼ Below Benchmark |
| | AMG | 66.0% | 66.4% | 60.0% ▼ Below Benchmark |
| | CCW | 71.8% | 69.7% ▼ | 67.5% ▼ At Benchmark |
| | CHPW | 67.0% | 64.3% ▼ | 61.0% ▼ Below Benchmark |
| | MHW | 67.8% | 66.4% ▼ | 64.6% ▼ Below Benchmark |
| | UHC | 63.6% | 63.5% | 61.1% ▼ Below Benchmark |
| Mental Health Treatment Penetration (MH-B), 6-26 years | Statewide† | 62.8% | 62.2% 🔻 | 60.4% ▼ Below Benchmark |
| | AMG | 60.0% | 62.0% 🔺 | 57.2% ▼ Below Benchmark |
| | CCW | 66.1% | 65.1% | 63.1% ▼ At Benchmark |
| | | | | |

| | CHPW MHW | 62.0% 63.2% | 60.5% ▼ 62.4% ▼ | 58.8% ▼ Below Benchmark 61.1% ▼ Below Benchmark |
|---|-------------|----------------|--------------------|---|
| | UHC | 58.4% | 58.9% | 57.6% Below Benchmark |
| Mental Health Treatment Penetration (MH-B), 18-64 years | Statewide† | 49.3% | 50.1% 🛕 | 50.1% At Benchmark |
| | AMG | 49.3% | 50.1% | 49.7% At Benchmark |
| | CCW | 48.4% | 49.2% | 48.3% Below Benchmark |
| | CHPW | 49.8% | 50.1% | 51.5% 🛕 Above Benchmark |
| | MHW | 50.1% | 51.2% 🔺 | 51.3% Above Benchmark |
| | UHC | 47.1% | 46.5% | 46.1% Below Benchmark |