

Comparison of Ideal vs. Actual Weight Base Factor Dosing

Background

Hypothesis: Factor dosing based on ideal body weight will result in hemostatic factor levels (recovery of at least 66% and a 6 hour half life)

Trial Design

This is a randomized, prospective, multicenter, open-label, cross-over study comparing the pharmacokinetics (PK) of ideal vs. actual body weight dosing of factor concentrate in patients with hemophilia.

The study will be conducted at the Washington Center for Bleeding Disorders (WCBD), Oregon Health & Science University (OHSU), Seattle Children's Hospital (SCH) and Sacred Heart Hospital (SH). Ethics approval will be obtained at all locations before trial enrollment.

Inclusion Criteria

- At least 2 years of age
- Hemophilia A or B – any severity
- Male gender
- Able and willing to consent to study
- Able and willing to comply with PK testing schedule
- Either overweight body weight (BMI 25 - 30) or obese (BMI > 30)

Exclusion Criteria

- Inhibitor of > 0.6 BU twice in the past
- Known other bleeding disorder
- Known other prolongation in aPTT (lupus anticoagulant, FXII deficiency)
- Female gender

Recruitment


Patients will be recruited through the participating centers.


Sample size

Do we want to differentiate between overweight and obese?

What should be the age cut-off (12 rather than 18)?


What is a feasible n?





Study Procedure:

How do we want to randomize?



PK protocol:

PK studies will be measure in response to one 100% corrective dose of the patient's current product. All patients will undergo PK testing twice: One with a 100% corrective dose (50U/kg for hemophilia A and 100U/kg for hemophilia B) base on ideal body weight and once based on actual body weight.

Ideal body weight will be calculated as follows:

$$50 + 2.3 (\text{height in inc.} - 60)$$

Hemophilia A – regular half-life product

Baseline – 30 min – 60 min – 2 hour – 4 hours – 6 hours – 8 hours – 24 hours – 48 hours

Hemophilia A – extended half-life factor

Baseline – 30 min – 60 min – 2 hour – 4 hours – 6 hours – 8 hours – 24 hours
– 48 hours – 72 hours – 96 hours

Hemophilia B – regular half-life product

Baseline – 30 min – 60 min – 2 hour – 4 hours – 6 hours – 8 hours – 24 hours
– 48 hours – 72 hours

Hemophilia B – extended half-life factor

Baseline – 30 min – 60 min – 2 hour – 4 hours – 6 hours – 8 hours – 24 hours
– 48 hours – 72 hours – 96 hours – 120 hours -