11/21/2023

Measure CCP: Contraceptive Care -- Postpartum Women Ages 15-20 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Three Days of Delivery By Year and Delivery System

		2018			2019			2020			2021			2022	
		Used Mo			Used Mo			Used Mo	•		Used Mo	•		Used Mo	*
		Effec			Effec			Effec			Effec			Effec	
		Contrac			Contrac			Contrac			Contrac			Contrac	-
	TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1	
	ELIGIBLE		% of												
Medicaid Managed Care Plan	WOMEN	N	Total												
Amerigroup Washington Inc	100	**	**	133	**	**	116	**	**	112	**	**	135	**	**
Community Health Plan of WA	234	**	**	223	**	**	186	**	**	170	**	**	188	**	**
Coordinated Care of WA	218	**	**	185	**	**	162	**	**	168	**	**	170	**	**
Molina Healthcare of WA Inc	771	15	1.9%	715	15	2.1%	726	15	2.1%	732	13	1.8%	635	15	2.4%
United Health Care Community Plan	92	**	**	73	**	**	80	**	**	87	**	**	71	**	**
Uncategorized	173	**	**	178	**	**	107	**	**	86	0	0.0%	75	**	**
Medicaid Managed Care	1,588	**	**	1,507	**	**	1,377	**	**	1,355	**	**	1,274	**	**
Medicaid Fee for Service	88	**	**	73	**	**	67	**	**	66	**	**	49	**	**
Total	1,676	38	2.3%	1,580	36	2.3%	1,444	33	2.3%	1,421	30	2.1%	1,323	29	2.2%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. **Excludes** women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 15-20 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Three Days of Delivery

By Year and Race/Ethnicity (Mutually Exclusive)

		2018			2019			2020			2021			2022	
		Used Mo	*		Used Mo										
		Effec			Effec			Effe			Effec			Effec	
		Contrac			Contrac			Contrac	·		Contrac			Contrac	
	TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1	15-20	TOTAL	Age 1		TOTAL	Age 1	
	ELIGIBLE		% of												
	WOMEN	N	Total												
Hispanic	614	12	2.0%	610	13	2.1%	622	16	2.6%	606	**	**	556	12	2.2%
Not Hispanic or Ethnicity Unknown															
White	693	13	1.9%	652	14	2.1%	542	**	**	515	**	**	476	11	2.3%
Asian	14	**	**	13	0	0.0%	12	0	0.0%	10	0	0.0%	18	**	**
Black	122	**	**	89	**	**	83	**	**	91	**	**	65	0	0.0%
American Indian/Alaska Native	106	**	**	90	**	**	78	**	**	86	**	**	67	**	**
Hawaiian/Pacific Islander	57	0	0.0%	55	0	0.0%	46	**	**	46	**	**	71	**	**
More Than One Race	28	0	0.0%	29	**	**	30	**	**	30	0	0.0%	23	0	0.0%
Other/Unknown	42	**	**	42	0	0.0%	31	0	0.0%	37	**	**	47	**	**
Total	1,676	38	2.3%	1,580	36	2.3%	1,444	33	2.3%	1,421	30	2.1%	1,323	29	2.2%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Race/ethnicity categories are mutually exclusive. Hispanic women may be of any race. **Rates** calculated from small numbers may be unstable and may fluctuate widely. Caution should be used when interpreting rates based on small numbers. ** = suppressed due to small numbers.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 15-20 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Three Days of Delivery

By Year and Race/Ethnicity (Not Mutually Exclusive)

		2018			2019			2020			2021			2022	
		Used Mo				re/Most			re/Most		Used Mo			Used Mo	
		Effec				ctive			ctive			ctive		Effec	
		Contrac				ception			ception			ception		Contrac	
	TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age .	L5-20 -	TOTAL	Age	L5-20	TOTAL	Age	L5-20 -
	ELIGIBLE		% of	ELIGIBLE		% of	ELIGIBLE		% of	ELIGIBLE		% of	ELIGIBLE		% of
	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total
American Indian/Alaska Native	137	**	**	121	**	**	118	**	**	117	**	**	98	**	**
Asian	22	**	**	18	0	0.0%	19	0	0.0%	15	0	0.0%	21	**	**
Black or African American	155	**	**	117	**	**	112	**	**	123	**	**	97	0	0.0%
Hispanic or Latino	614	12	2.0%	610	13	2.1%	622	16	2.6%	606	**	**	556	12	2.2%
Native Hawaiian/Pacific Islander	69	0	0.0%	75	**	**	64	**	**	66	**	**	83	**	**
White, Non-Hispanic (and none of the above)	693	13	1.9%	652	14	2.1%	542	**	**	515	**	**	476	11	2.3%
Other/Unknown	42	**	**	42	0	0.0%	31	0	0.0%	37	**	**	47	**	**
Total	1,676	38	2.3%	1,580	36	2.3%	1,444	33	2.3%	1,421	30	2.1%	1,323	29	2.2%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Race/ethnicity categories are not mutually exclusive with the exception of White, Non-Hispanic, which means women with multiple race/ethnicity categories are counted in each applicable category. Rates calculated from small numbers may be unstable and may fluctuate widely. Caution should be used when interpreting rates based on small numbers. ** = suppressed due to small numbers.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 21-44 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Three Days of Delivery By Year and Delivery System

		2018			2019			2020			2021			2022	
		Used Mo	1		Used Mo			Used Mo			Used Mo				re/Most
		Effec			Effec			Effec			Effec				ctive
		Contrac			Contrac			Contrac			Contrac				ception
	TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2	21-44
	ELIGIBLE		% of	ELIGIBLE		% of									
Medicaid Managed Care Plan	WOMEN	N	Total	WOMEN	N	Total									
Amerigroup Washington Inc	1,333	147	11.0%	1,474	119	8.1%	1,738	159	9.1%	1,707	150	8.8%	1,762	139	7.9%
Community Health Plan of WA	2,185	201	9.2%	1,903	160	8.4%	1,634	162	9.9%	1,714	148	8.6%	1,861	141	7.6%
Coordinated Care of WA	1,635	169	10.3%	1,351	148	11.0%	1,465	124	8.5%	1,453	127	8.7%	1,561	150	9.6%
Molina Healthcare of WA Inc	7,602	805	10.6%	7,369	709	9.6%	8,707	853	9.8%	9,110	769	8.4%	9,316	770	8.3%
United Health Care Community Plan	1,761	151	8.6%	1,234	89	7.2%	1,637	134	8.2%	1,652	140	8.5%	1,643	109	6.6%
Uncategorized	3,197	240	7.5%	3,030	227	7.5%	1,132	92	8.1%	843	42	5.0%	750	44	5.9%
Medicaid Managed Care	17,713	1,713	9.7%	16,361	1,452	8.9%	16,313	1,524	9.3%	16,479	1,376	8.4%	16,893	1,353	8.0%
Medicaid Fee for Service	603	79	13.1%	482	51	10.6%	428	43	10.0%	369	40	10.8%	333	27	8.1%
Total	18,316	1,792	9.8%	16,843	1,503	8.9%	16,741	1,567	9.4%	16,848	1,416	8.4%	17,226	1,380	8.0%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 21-44 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Three Days of Delivery

By Year and Race/Ethnicity (Mutually Exclusive)

		2018			2019			2020			2021			2022	
		Used Mo	,			re/Most		Used Mo	•		Used Mo			Used Mo	*
		Effec			Effec			Effec			Effec			Effec	
		Contrac			Contrac	•		Contrac	•		Contrac			Contrac	•
	TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2	
	ELIGIBLE	NI	% of	ELIGIBLE	NI.	% of	ELIGIBLE	NI	% of	ELIGIBLE	N.	% of	ELIGIBLE	NI.	% of
	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total
Hispanic	3,869	395	10.2%	3,778	366	9.7%	3,856	365	9.5%	3,991	364	9.1%	4,333	389	9.0%
Not Hispanic or Ethnicity Unknown															
White	9,751	985	10.1%	8,683	825	9.5%	8,488	834	9.8%	8,459	720	8.5%	8,125	671	8.3%
Asian	657	30	4.6%	582	32	5.5%	575	31	5.4%	582	34	5.8%	644	41	6.4%
Black	1,570	144	9.2%	1,454	96	6.6%	1,495	103	6.9%	1,504	98	6.5%	1,556	102	6.6%
American Indian/Alaska Native	777	104	13.4%	659	67	10.2%	665	76	11.4%	671	71	10.6%	642	47	7.3%
Hawaiian/Pacific Islander	742	62	8.4%	744	58	7.8%	791	81	10.2%	772	68	8.8%	925	67	7.2%
More Than One Race	203	18	8.9%	229	15	6.6%	197	24	12.2%	216	22	10.2%	277	23	8.3%
Other/Unknown	747	54	7.2%	714	44	6.2%	674	53	7.9%	653	39	6.0%	724	40	5.5%
Total	18,316	1,792	9.8%	16,843	1,503	8.9%	16,741	1,567	9.4%	16,848	1,416	8.4%	17,226	1,380	8.0%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Race/ethnicity categories are mutually exclusive. Hispanic women may be of any race. Rates calculated from small numbers may be unstable and may fluctuate widely. Caution should be used when interpreting rates based on small numbers. ** = suppressed due to small numbers or for secondary suppression.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 21-44 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Three Days of Delivery

By Year and Race/Ethnicity (Not Mutually Exclusive)

		2018			2019			2020			2021			2022	
		Used Mo	1		Used Mo			Used Mo	•		Used Mo	•		Used Mo	*
		Effec			Effec				ctive		Effec			Effec	
	TOTAL	Contrac		TOTAL	Contrac	•	TOT41		ception	TOTAL	Contrac	·	TOTAL	Contrac	
	TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2	
	ELIGIBLE	N	% of	ELIGIBLE	N	% of	ELIGIBLE	NI	% of	ELIGIBLE	NI	% of	ELIGIBLE	N	% of
	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total
American Indian/Alaska Native	969	127	13.1%	834	86	10.3%	836	90	10.8%	850	86	10.1%	815	57	7.0%
Asian	729	34	4.7%	665	36	5.4%	650	36	5.5%	666	41	6.2%	749	52	6.9%
Black or African American	1,763	161	9.1%	1,665	114	6.8%	1,707	128	7.5%	1,723	122	7.1%	1,843	130	7.1%
Hispanic or Latino	3,869	395	10.2%	3,778	366	9.7%	3,856	365	9.5%	3,991	364	9.1%	4,333	389	9.0%
Native Hawaiian/Pacific Islander	866	73	8.4%	862	65	7.5%	914	94	10.3%	904	83	9.2%	1,078	79	7.3%
White, Non-Hispanic (and none of the above)	9,751	985	10.1%	8,683	825	9.5%	8,488	834	9.8%	8,459	720	8.5%	8,125	671	8.3%
Other/Unknown	747	54	7.2%	714	44	6.2%	674	53	7.9%	653	39	6.0%	724	40	5.5%
Total	18,316	1,792	9.8%	16,843	1,503	8.9%	16,741	1,567	9.4%	16,848	1,416	8.4%	17,226	1,380	8.0%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Race/ethnicity categories are not mutually exclusive with the exception of White, Non-Hispanic, which means women with multiple race/ethnicity categories are counted in each applicable category. Rates calculated from small numbers may be unstable and may fluctuate widely. Caution should be used when interpreting rates based on small numbers. ** = suppressed due to small numbers or for secondary suppression.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 15-20 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Ninety Days of Delivery By Year and Delivery System

		2018			2019			2020			2021			2022	
		Used Mo	•		Used Mo			Used Mo	•		Used Mo				ore/Most
		Effec			Effec			Effec			Effec				ctive
		Contrac			Contrac			Contrac	·		Contrac	•			ception
	TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1	L5-20
	ELIGIBLE		% of	ELIGIBLE		% of									
Medicaid Managed Care Plan	WOMEN	N	Total	WOMEN	N	Total									
Amerigroup Washington Inc	100	56	56.0%	133	69	51.9%	116	62	53.4%	112	63	56.3%	135	67	49.6%
Community Health Plan of WA	234	125	53.4%	223	132	59.2%	186	111	59.7%	170	83	48.8%	188	99	52.7%
Coordinated Care of WA	218	123	56.4%	185	113	61.1%	162	102	63.0%	168	82	48.8%	170	92	54.1%
Molina Healthcare of WA Inc	771	449	58.2%	715	437	61.1%	726	404	55.6%	732	373	51.0%	635	288	45.4%
United Health Care Community Plan	92	54	58.7%	73	40	54.8%	80	51	63.8%	87	39	44.8%	71	25	35.2%
Uncategorized	173	87	50.3%	178	94	52.8%	107	60	56.1%	86	39	45.3%	75	28	37.3%
Medicaid Managed Care	1,588	894	56.3%	1,507	885	58.7%	1,377	790	57.4%	1,355	679	50.1%	1,274	599	47.0%
Medicaid Fee for Service	88	39	44.3%	73	38	52.1%	67	35	52.2%	66	27	40.9%	49	22	44.9%
Total	1,676	933	55.7%	1,580	923	58.4%	1,444	825	57.1%	1,421	706	49.7%	1,323	621	46.9%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. **Excludes** women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 15-20 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Ninety Days of Delivery

By Year and Race/Ethnicity (Mutually Exclusive)

		2018			2019			2020			2021			2022	
		Used Mo			Used Mo	•		Used Mo			Used Mo			Used Mo	
		Effec			Effec			Effec			Effec			Effec	
		Contrac			Contrac			Contrac	•		Contrac			Contrac	
	TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age 1	
	ELIGIBLE		% of												
	WOMEN	N	Total												
Hispanic	614	351	57.2%	610	375	61.5%	622	392	63.0%	606	330	54.5%	556	293	52.7%
Not Hispanic or Ethnicity Unknown															
White	693	407	58.7%	652	399	61.2%	542	292	53.9%	515	254	49.3%	476	221	46.4%
Asian	14	**	**	13	**	**	12	**	**	10	**	**	18	**	**
Black	122	54	44.3%	89	44	49.4%	83	40	48.2%	91	39	42.9%	65	20	30.8%
American Indian/Alaska Native	106	52	49.1%	90	45	50.0%	78	44	56.4%	86	35	40.7%	67	29	43.3%
Hawaiian/Pacific Islander	57	23	40.4%	55	19	34.5%	46	17	37.0%	46	**	**	71	25	35.2%
More Than One Race	28	19	67.9%	29	14	48.3%	30	19	63.3%	30	15	50.0%	23	**	**
Other/Unknown	42	**	**	42	**	**	31	**	**	37	18	48.6%	47	19	40.4%
Total	1,676	933	55.7%	1,580	923	58.4%	1,444	825	57.1%	1,421	706	49.7%	1,323	621	46.9%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Race/ethnicity categories are mutually exclusive. Hispanic women may be of any race. Rates calculated from small numbers may be unstable and may fluctuate widely. Caution should be used when interpreting rates based on small numbers. ** = suppressed due to small numbers or for secondary suppression.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 15-20 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Ninety Days of Delivery

By Year and Race/Ethnicity (Not Mutually Exclusive)

		2018			2019			2020			2021			2022	
		Used Mo	•		Used Mo				re/Most		Used Mo	•		Used Mo	*
		Effec				ctive			ctive			ctive			ctive
	TOTAL		ception	TOTAL	Contrac	•	TOTAL		ception	TOTAL		ception	TOTAL		ception
	TOTAL	Age 1		TOTAL	Age 1		TOTAL	Age	15-20	TOTAL	Age 1		TOTAL	Age	15-20
	ELIGIBLE	NI	% of	ELIGIBLE	NI.	% of	ELIGIBLE	NI.	% of	ELIGIBLE	N	% of	ELIGIBLE	NI.	% of
	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total	WOMEN	N	Total
American Indian/Alaska Native	137	69	50.4%	121	60	49.6%	118	66	55.9%	117	50	42.7%	98	41	41.8%
Asian	22	**	**	18	**	**	19	11	57.9%	15	**	**	21	**	**
Black or African American	155	74	47.7%	117	57	48.7%	112	58	51.8%	123	58	47.2%	97	31	32.0%
Hispanic or Latino	614	351	57.2%	610	375	61.5%	622	392	63.0%	606	330	54.5%	556	293	52.7%
Native Hawaiian/Pacific Islander	69	31	44.9%	75	33	44.0%	64	29	45.3%	66	19	28.8%	83	31	37.3%
White, Non-Hispanic (and none of the above)	693	407	58.7%	652	399	61.2%	542	292	53.9%	515	254	49.3%	476	221	46.4%
Other/Unknown	42	**	**	42	**	**	31	14	45.2%	37	**	**	47	**	**
Total	1,676	933	55.7%	1,580	923	58.4%	1,444	825	57.1%	1,421	706	49.7%	1,323	621	46.9%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Race/ethnicity categories are not mutually exclusive with the exception of White, Non-Hispanic, which means women with multiple race/ethnicity categories are counted in each applicable category. Rates calculated from small numbers may be unstable and may fluctuate widely. Caution should be used when interpreting rates based on small numbers.

** = suppressed due to small numbers or for secondary suppression.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 21-44 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Ninety Days of Delivery By Year and Delivery System

		2018			2019			2020			2021			2022	
		Used Mo			Used Mo			Used Mo	•		Used Mo	•		Used Mo	
		Effec			Effec			Effec			Effec			Effec	
		Contrac			Contrac			Contrac	·		Contrac			Contrac	•
	TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2	
	ELIGIBLE		% of												
Medicaid Managed Care Plan	WOMEN	N	Total												
Amerigroup Washington Inc	1,333	672	50.4%	1,474	681	46.2%	1,738	805	46.3%	1,707	731	42.8%	1,762	713	40.5%
Community Health Plan of WA	2,185	1,074	49.2%	1,903	984	51.7%	1,634	781	47.8%	1,714	816	47.6%	1,861	849	45.6%
Coordinated Care of WA	1,635	934	57.1%	1,351	767	56.8%	1,465	743	50.7%	1,453	720	49.6%	1,561	712	45.6%
Molina Healthcare of WA Inc	7,602	3,859	50.8%	7,369	3,708	50.3%	8,707	4,123	47.4%	9,110	3,996	43.9%	9,316	3,762	40.4%
United Health Care Community Plan	1,761	848	48.2%	1,234	584	47.3%	1,637	736	45.0%	1,652	704	42.6%	1,643	589	35.8%
Uncategorized	3,197	1,368	42.8%	3,030	1,368	45.1%	1,132	500	44.2%	843	294	34.9%	750	240	32.0%
Medicaid Managed Care	17,713	8,755	49.4%	16,361	8,092	49.5%	16,313	7,688	47.1%	16,479	7,261	44.1%	16,893	6,865	40.6%
Medicaid Fee for Service	603	270	44.8%	482	215	44.6%	428	168	39.3%	369	159	43.1%	333	109	32.7%
Total	18,316	9,025	49.3%	16,843	8,307	49.3%	16,741	7,856	46.9%	16,848	7,420	44.0%	17,226	6,974	40.5%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. **Excludes** women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 21-44 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Ninety Days of Delivery

By Year and Race/Ethnicity (Mutually Exclusive)

		2018			2019			2020			2021			2022	
		Used Mo			Used Mo			Used Mo			Used Mo			Used Mo	*
		Effec			Effec			Effec			Effec			Effec	
		Contrac			Contrac			Contrac			Contrac			Contrac	
	TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2	
	ELIGIBLE		% of												
	WOMEN	N	Total												
Hispanic	3,869	2,248	58.1%	3,778	2,206	58.4%	3,856	2,112	54.8%	3,991	2,135	53.5%	4,333	2,133	49.2%
Not Hispanic or Ethnicity Unknown															
White	9,751	4,701	48.2%	8,683	4,223	48.6%	8,488	3,900	45.9%	8,459	3,571	42.2%	8,125	3,043	37.5%
Asian	657	260	39.6%	582	237	40.7%	575	238	41.4%	582	236	40.5%	644	229	35.6%
Black	1,570	739	47.1%	1,454	613	42.2%	1,495	630	42.1%	1,504	571	38.0%	1,556	613	39.4%
American Indian/Alaska Native	777	358	46.1%	659	317	48.1%	665	291	43.8%	671	300	44.7%	642	234	36.4%
Hawaiian/Pacific Islander	742	309	41.6%	744	304	40.9%	791	329	41.6%	772	278	36.0%	925	323	34.9%
More Than One Race	203	105	51.7%	229	96	41.9%	197	88	44.7%	216	89	41.2%	277	115	41.5%
Other/Unknown	747	305	40.8%	714	311	43.6%	674	268	39.8%	653	240	36.8%	724	284	39.2%
Total	18,316	9,025	49.3%	16,843	8,307	49.3%	16,741	7,856	46.9%	16,848	7,420	44.0%	17,226	6,974	40.5%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Race/ethnicity categories are mutually exclusive. Hispanic women may be of any race. **Rates** calculated from small numbers may be unstable and may fluctuate widely. Caution should be used when interpreting rates based on small numbers. ** = suppressed due to small numbers.

Measure CCP: Contraceptive Care -- Postpartum Women Ages 21-44 Percentage Who Were Provided a Most Effective or Moderately Effective FDA-approved Method of Contraception Within Ninety Days of Delivery

By Year and Race/Ethnicity (Not Mutually Exclusive)

		2018			2019			2020			2021			2022	
		Used Mo			Used Mo	•		Used Mo			Used Mo	•		Used Mo	*
		Effec Contrac			Effect Contract			Effec			Effec	ctive ception		Effec	ctive ception
	TOTAL	Age 2		TOTAL	Age 2	·	TOTAL	Contrac Age 2		TOTAL	Age 2	·	TOTAL		21-44
	TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2		TOTAL	Age 2	
	ELIGIBLE WOMEN	N	% of Total	ELIGIBLE WOMEN	N	% of Total	ELIGIBLE WOMEN	N	% of Total	ELIGIBLE WOMEN	N	% of Total	ELIGIBLE WOMEN	N	% of Total
American Indian/Alaska Native	969		46.9%	834		47.5%			43.2%	850		44.4%			36.1%
•															
Asian	729	298	40.9%	665	276	41.5%	650	279	42.9%	666	269	40.4%	749	273	36.4%
Black or African American	1,763	821	46.6%	1,665	704	42.3%	1,707	723	42.4%	1,723	653	37.9%	1,843	729	39.6%
Hispanic or Latino	3,869	2,248	58.1%	3,778	2,206	58.4%	3,856	2,112	54.8%	3,991	2,135	53.5%	4,333	2,133	49.2%
Native Hawaiian/Pacific Islander	866	378	43.6%	862	359	41.6%	914	384	42.0%	904	332	36.7%	1,078	386	35.8%
White, Non-Hispanic (and none of the above)	9,751	4,701	48.2%	8,683	4,223	48.6%	8,488	3,900	45.9%	8,459	3,571	42.2%	8,125	3,043	37.5%
Other/Unknown	747	305	40.8%	714	311	43.6%	674	268	39.8%	653	240	36.8%	724	284	39.2%
Total	18,316	9,025	49.3%	16,843	8,307	49.3%	16,741	7,856	46.9%	16,848	7,420	44.0%	17,226	6,974	40.5%

Most or moderately effective FDA-approved contraception methods: female sterilization, contraceptive implants, intrauterine devices or systems, injectables, oral pills, patch, or ring.

Eligible women had a live birth in the first nine months of the measurement year, were continuously enrolled in Medicaid or CHIP with medical or family planning benefits from the date of delivery to 90 days postpartum, and were in the specified age range as of December 31 of the measurement year. Excludes women who were eligible for both Medicaid and Medicare; women who had full third party liability; women with deliveries that did not end in a live birth.

Race/ethnicity categories are not mutually exclusive with the exception of White, Non-Hispanic, which means women with multiple race/ethnicity categories are counted in each applicable category. Rates calculated from small numbers may be unstable and may fluctuate widely. Caution should be used when interpreting rates based on small numbers.

** = suppressed due to small numbers.