

Chlamydia Screening in Women

Metric Information

Metric description: The percentage of female Medicaid beneficiaries, 16–24 years of age identified as sexually active and who had at least one test for chlamydia during the measurement year.

Metric specification version: HEDIS® Measurement Year 2020 and Measurement Year 2021 Technical Specifications for Health Plans, NCQA.

Data collection method: Administrative only.

Data source: ProviderOne Medicaid claims/encounter and enrollment data.

Claim status: Include only final paid claims or accepted encounters in metric calculation.

Identification window: Measurement year.

Direction of quality improvement: Higher is better.

URL of specifications: www.ncqa.org/hedis/measures

DSRIP Program Summary

Metric utility: ACH Project P4P ACH High Performance DSRIP statewide accountability

ACH Project P4P – Metric results used for achievement value: Single metric result.

ACH Project P4P – improvement target methodology: gap to goal.

ACH Project P4P gap to goal - absolute benchmark value:

DY 3/performance year 1 (2019)	71.5% 2017 NCQA Quality Compass National Medicaid, 90 th Percentile
DY 4/performance year 2 (2020)	71.33% 2018 NCQA Quality Compass National Medicaid, 90 th Percentile
DY 5/performance year 3 (2021)	71.42% 2019 NCQA Quality Compass National Medicaid, 90 th Percentile

ACH regional attribution: Residence in the ACH region for 11 out of 12 months in the measurement year.

DSRIP Metric Details

Eligible Population	
Age	16-24 years. Age is as of the last day of the measurement year.
Gender	Female

Chlamydia Screening in Women

Minimum Medicaid enrollment	Measurement year. Continuous enrollment is required.
Allowable gap in Medicaid enrollment	One gap of one month during the measurement year.
Medicaid enrollment anchor date	Last day of measurement year.
Medicaid benefit and eligibility	Includes Medicaid beneficiaries with comprehensive medical benefits or family planning only benefits. Excludes beneficiaries that are eligible for both Medicare and Medicaid and beneficiaries with primary insurance other than Medicaid.

Denominator:

Data elements required for denominator: Sexually active females need to be identified for this metric. There are two methods to identify sexually active women: claim/encounter data and pharmacy data. Both methods are used to identify eligible female Medicaid beneficiaries; however, a beneficiary only needs to be identified in one method to be eligible for this metric.

Required exclusions for denominator.

- Eligible population exclusions are listed in the eligible population table above.
- Metric specific exclusions:
 - o Beneficiaries in hospice care.
 - o Excludes women identified as sexually active solely due to a pregnancy test and who had an x-ray or prescription for isotretinoin within seven days of the pregnancy test.

Deviations from cited specifications for denominator.

- HEDIS® specifications require no more than one gap in continuous enrollment of up to 45 days during each year of continuous enrollment. To determine continuous enrollment for a Medicaid beneficiary for whom enrollment is verified monthly, as is the case for the ProviderOne data source, the Medicaid beneficiary may not have more than a 1-month gap in coverage (i.e., a Medicaid beneficiary whose coverage lapses for 2 months [60 days] is not considered continuously enrolled).

Numerator:

Beneficiaries must qualify for inclusion in the denominator to be eligible for inclusion in the numerator.

Data elements required for numerator. At least one chlamydia test during the measurement year.

Required exclusions for numerator.

- None

Deviations from cited specifications for numerator.

- None

Chlamydia Screening in Women

Version Control